

Colouring Square-Free Graphs without Long Induced Paths

Serge Gaspers

UNSW Sydney and Data61, CSIRO, Australia
sergeg@cse.unsw.edu.au

Shenwei Huang

UNSW Sydney, Australia
dynamichuang@gmail.com

Daniël Paulusma

Durham University, Durham, UK
daniel.paulusma@durham.ac.uk

Abstract

The COLOURING problem is to decide if the vertices of a graph can be coloured with at most k colours for a given integer k such that no two adjacent vertices are coloured alike. The complexity of COLOURING is fully understood for graph classes characterized by one forbidden induced subgraph H . Despite a huge body of existing work, there are still major complexity gaps if two induced subgraphs H_1 and H_2 are forbidden. We let H_1 be the s -vertex cycle C_s and H_2 be the t -vertex path P_t . We show that COLOURING is polynomial-time solvable for $s = 4$ and $t \leq 6$, which unifies several known results for COLOURING on (H_1, H_2) -free graphs. Our algorithm is based on a novel decomposition theorem for (C_4, P_6) -free graphs without clique cutsets into homogeneous pairs of sets and a new framework for bounding the clique-width of a graph by the clique-width of its subgraphs induced by homogeneous pairs of sets. To apply this framework, we also need to use divide-and-conquer to bound the clique-width of subgraphs induced by homogeneous pairs of sets. To complement our positive result we also prove that COLOURING is NP-complete for $s = 4$ and $t \geq 9$, which is the first hardness result on COLOURING for (C_4, P_t) -free graphs.

2012 ACM Subject Classification Mathematics of computing → Graph theory

Keywords and phrases Graph Colouring, Hereditary Graph Class, Clique-width, Cycle, Path

Digital Object Identifier 10.4230/LIPIcs.STACS.2018.35

Funding Serge Gaspers is the recipient of an Australian Research Council (ARC) Future Fellowship (FT140100048) and acknowledges support under the ARC's Discovery Projects funding scheme (DP150101134). Daniël Paulusma is supported by Leverhulme Trust Grant RPG-2016-258.

Acknowledgements Initially we proved NP-hardness of COLOURING for (C_4, P_{16}) -free graphs. Afterwards we were able to improve this result to (C_4, P_9) -free graphs via a simplification of our construction. We would like to thank an anonymous reviewer for pointing out this simplification as well.



© Serge Gaspers, Shenwei Huang, and Daniël Paulusma;
licensed under Creative Commons License CC-BY
35th Symposium on Theoretical Aspects of Computer Science (STACS 2018).
Editors: Rolf Niedermeier and Brigitte Vallée; Article No. 35; pp. 35:1–35:15
Leibniz International Proceedings in Informatics



LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany



SYMPOSIUM
ON THEORETICAL
ASPECTS
OF COMPUTER
SCIENCE

1 Introduction

Graph colouring has been a popular and extensively studied concept in computer science and mathematics since its introduction as a map colouring problem more than 150 years ago due to its many application areas crossing disciplinary boundaries and to its use as a benchmark problem in research into computational hardness. The corresponding decision problem, COLOURING, is to decide, for a given graph G and integer k , if G admits a k -colouring, that is, a mapping $c : V(G) \rightarrow \{1, \dots, k\}$ such that $c(u) \neq c(v)$ whenever $uv \in E(G)$. Unless $P = NP$, it is not possible to solve COLOURING in polynomial time for general graphs, not even if the number of colours is limited to 3 [37]. To get a better understanding of the borderline between tractable and intractable instances of COLOURING, it is natural to restrict the input to some special graph class. Hereditary graph classes, which are classes of graphs closed under vertex deletion, provide a unified framework for a large collection of well-known graph classes. It is readily seen that a graph class is hereditary if and only if it can be characterized by a (unique) set \mathcal{H} of minimal forbidden induced subgraphs. Graphs with no induced subgraph isomorphic to a graph in a set \mathcal{H} are called \mathcal{H} -free.

Over the years, the study of COLOURING for hereditary graph classes has evolved into a deep area of research in theoretical computer science and discrete mathematics (see, for example, [6, 22, 31, 44]). One of the best-known results is the classical result of Grötschel, Lovász, and Schrijver [24], who showed that COLOURING is polynomial-time solvable for perfect graphs. Faster, even linear-time, algorithms are known for subclasses of perfect graphs, such as chordal graphs, bipartite graphs, interval graphs, and comparability graphs; see for example [22]. All these classes are characterized by infinitely many minimal forbidden induced subgraphs. Král', Kratochvíl, Tuza, and Woeginger [35] initiated a systematic study into the computational complexity of COLOURING restricted to hereditary graph classes characterized by a *finite* number of minimal forbidden induced subgraphs. In particular they gave a complete classification of the complexity of COLOURING for the case where \mathcal{H} consists of a single graph H . Their dichotomy result led to two natural directions for further research:

1. Is it possible to obtain a dichotomy for COLOURING on H -free graphs if the number of colours, k , is fixed (that is, k no longer belongs to the input)?
2. Is it possible to obtain a dichotomy for COLOURING on \mathcal{H} -free graphs if \mathcal{H} has size 2?

We briefly discuss known results for both directions below and refer to [19] for a detailed survey. Let C_s and P_t denote the cycle on s vertices and path on t vertices, respectively. We start with the first question. If k is fixed, then we denote the problem by k -COLOURING. It is known that for every $k \geq 3$, the k -COLOURING problem on H -free graphs is NP-complete whenever H contains a cycle [16] or an induced claw [28, 36]. Therefore, only the case when H is a disjoint union of paths remains. In particular, the situation where $H = P_t$ has been thoroughly studied. On the positive side, 3-COLOURING P_7 -free graphs and k -COLOURING P_5 -free graphs for any fixed $k \geq 1$ are shown to be polynomial-time solvable [3, 26]. On the negative side, Huang [29] proved NP-completeness for $(k = 5, t = 6)$ and for $(k = 4, t = 7)$. The cases $(k = 3, t \geq 8)$ and $(k = 4, t = 6)$ remain open, although some partial results are known [9, 10].

In this paper we focus on the second question, that is, we restrict the input of COLOURING to \mathcal{H} -free graphs for $\mathcal{H} = \{H_1, H_2\}$. For two graphs G and H , we use $G + H$ to denote the disjoint union of G and H , and we write rG to denote the disjoint union of r copies of G . As a starting point, Král', Kratochvíl, Tuza, and Woeginger [35] identified the following three main sources of NP-completeness: (i) both H_1 and H_2 contain a claw; (ii) both H_1 and H_2 contain a cycle; and (iii) both H_1 and H_2 contain an induced subgraph from the set

$\{4P_1, 2P_1 + P_2, 2P_2\}$. They also showed additional NP-completeness results by mixing the three types. Since then numerous papers [1, 7, 8, 13, 14, 25, 27, 29, 33, 35, 38, 41, 42, 43, 47] have been devoted to this problem, but despite all these efforts the complexity classification for COLOURING on (H_1, H_2) -free graphs is still far from complete, and even dealing with specific pairs (H_1, H_2) may require substantial work.

One of the “mixed” results obtained in [35] is that COLOURING is NP-complete for (C_s, H) -free graphs when $s \geq 5$ and $H \in \{4P_1, 2P_1 + P_2, 2P_2\}$. This, together with the well-known result that COLOURING can be solved in linear time for P_4 -free graphs, implies the following dichotomy.

► **Theorem 1** ([35]). *Let $s \geq 5$ be a fixed positive integer. Then COLOURING for (C_s, P_t) -free graphs is polynomial-time solvable when $t \leq 4$ and NP-complete when $t \geq 5$.*

Theorem 1 raises the natural question: what is the complexity of COLOURING on (C_s, P_t) -free graphs when $s \in \{3, 4\}$? Huang, Johnson and Paulusma [30] proved that 4-COLOURING, and thus COLOURING, is NP-complete for (C_3, P_{22}) -free graphs, while a result of Brandstädt, Klemmt and Mahfud [5] implies that COLOURING is polynomial-time solvable for (C_3, P_6) -free graphs. For $s = 4$, it is only known that COLOURING is polynomial-time solvable for (C_4, P_5) -free graphs [41]. This is unless we fix the number of colours: for every $k \geq 1$ and $t \geq 1$, it is known that k -COLOURING is polynomial-time solvable for (C_4, P_t) -free graphs [21].

Our Results. We first show, in section 3, that COLOURING is polynomial-time solvable for (C_4, P_6) -free graphs. This case was explicitly mentioned as a natural case to consider in [19]. Our result unifies several previous results on colouring (C_4, P_t) -free graphs, namely: the polynomial-time solvability of COLOURING for (C_4, P_5) -free graphs [41]; the polynomial-time solvability of k -COLOURING for (C_4, P_6) -free graphs for every $k \geq 1$ [21]; and the recent 3/2-approximation algorithm for COLOURING for (C_4, P_6) -free graphs [18]. It was not previously known if there exists an integer t such that COLOURING is NP-complete for (C_4, P_t) -free graphs. In section 4 we complement our positive result by giving an affirmative answer to this question: already the value $t = 9$ makes the problem NP-complete.

Our Methodology. The general research aim of our paper is to increase, in a systematic way, our insights in the computational hardness of COLOURING and to narrow the complexity gaps between hard and easy cases. Clique-width is a well-known width parameter and having bounded clique-width is often the underlying reason for a large collection of NP-complete problems, including COLOURING, to become tractable on a special graph class; this follows from results of [11, 17, 34, 45, 46]. However, the class of (C_4, P_6) -free graphs contains the class of split graphs, which may have arbitrarily large clique-width [40]. Hence, if we want to use clique-width to solve COLOURING for (C_4, P_6) -free graphs, then we first need to preprocess the input graph. An *atom* is a graph with no clique cut set. In this paper we prove that (C_4, P_6) -free atoms *have* bounded clique-width. This implies a polynomial-time algorithm for COLOURING on (C_4, P_6) -free graphs, as it is well known that COLOURING is polynomial-time solvable on a hereditary graph class \mathcal{G} if it is so on the atoms of \mathcal{G} [49].

In order to prove that (C_4, P_6) -free atoms have bounded clique-width, we further develop the approach of [18] that was used to bound the chromatic number of (C_4, P_6) -free graphs as a linear function of their maximum clique size and to obtain a 3/2-approximation algorithm for COLOURING for (C_4, P_6) -free graphs. The approach of [18] is based on a decomposition theorem for (C_4, P_6) -free atoms. For our purposes we derive a new variant of this decomposition theorem for so-called strong atoms, which are atoms that contain no universal

vertices and no pairs of twin vertices. Another novel element in our approach is that we show how to bound the clique-width of a graph by the clique-width of its subgraphs induced by homogeneous pairs of sets, and this will be very useful for dealing with (C_4, P_6) -free strong atoms. To apply this method, we also need to use divide-and-conquer to bound the clique-width of subgraphs induced by homogeneous pairs of sets.

2 Preliminaries

For general graph theory notation we follow [2]. Let $G = (V, E)$ be a graph. The *neighbourhood* of a vertex v , denoted by $N_G(v)$, is the set of neighbours of v . For a set $X \subseteq V(G)$, let $N_G(X) = \bigcup_{v \in X} N_G(v) \setminus X$. The *degree* of v , denoted by $d_G(v)$, is equal to $|N_G(v)|$. For $x \in V$ and $S \subseteq V$, we denote by $N_S(x)$ the set of neighbours of x that are in S , i.e., $N_S(x) = N_G(x) \cap S$. For $X, Y \subseteq V$, we say that X is *complete* (resp. *anti-complete*) to Y if every vertex in X is adjacent (resp. non-adjacent) to every vertex in Y . A vertex subset $K \subseteq V$ is a *clique cutset* if $G - K$ has more components than G and K induces a clique. A vertex is *universal* in G if it is adjacent to all other vertices. For $S \subseteq V$, the subgraph induced by S , is denoted by $G[S]$.

A subset $D \subseteq V$ is a *dominating set* if every vertex not in D has a neighbour in D . Let $u, v \in V$ be two distinct vertices. We say that a vertex $x \notin \{u, v\}$ *distinguishes* u and v if x is adjacent to exactly one of u and v . A set $H \subseteq V$ is a *homogeneous set* if no vertex in $V \setminus H$ can distinguish two vertices in H . A homogeneous set H is *proper* if $1 < |H| < |V|$. A graph is *prime* if it contains no proper homogeneous set. We say that u and v are *twins* if u and v are adjacent and they have the same set of neighbours in $V \setminus \{u, v\}$. Note that the binary relation of being twins is an equivalence relation on V and so V can be partitioned into equivalence classes T_1, \dots, T_r of twins. The *skeleton* of G is the subgraph induced by a set of r vertices, one from each of T_1, \dots, T_r . A *blow-up* of a graph G is a graph G' obtained by replacing each vertex v of G with a clique K_v of size at least 1 such that K_v and K_u are complete in G' if u and v are adjacent in G , and anti-complete otherwise. Since each equivalence class of twins is a clique and any two equivalence classes are either complete or anti-complete, every graph is a blow-up of its skeleton.

The *clique-width* of a graph G , denoted by $cw(G)$, is the minimum number of labels required to construct G using the following four operations:

- $i(v)$: create a new graph consisting of a single vertex v with label i ;
- $G_1 \oplus G_2$: take the disjoint union of two labelled graphs G_1 and G_2 ;
- $\eta_{i,j}$: join each vertex with label i to each vertex with label j (for $i \neq j$);
- $\rho_{i \rightarrow j}$: rename label i to j .

A *clique-width expression* for G is an algebraic expression that describes how G can be recursively constructed using these operations. A *k-expression* for G is a clique-width expression using at most k distinct labels. For instance, this is a 3-expression for the induced path on four vertices a, b, c, d :

$$\eta_{3,2}(3(d) \oplus \rho_{3 \rightarrow 2}(\rho_{2 \rightarrow 1}(\eta_{3,2}(3(c) \oplus \eta_{2,1}(2(b) \oplus 1(a)))))).$$

Clique-width is of fundamental importance in computer science since all problems expressible in monadic second-order logic using quantifiers over vertex subsets but not over edge subsets become polynomial-time solvable for graphs of bounded clique-width [11]. Although this meta-theorem does not directly apply to COLOURING, a result of Kobler and Rotics [34], combined with the approximation algorithm of Oum and Seymour [45] for finding a p -expression, showed that COLOURING can be added to the list of such problems.

► **Theorem 2** ([34]). COLOURING can be solved in polynomial time for graphs of bounded clique-width.

3 The Polynomial-Time Result

In this section, we shall prove that the chromatic number of any (C_4, P_6) -free graph can be found in polynomial time.

► **Theorem 3.** COLOURING is polynomial-time solvable on the class of (C_4, P_6) -free graphs.

A graph is called an *atom* if it contains no clique cutset. The main ingredient for proving Theorem 3 is a new structural property of (C_4, P_6) -free atoms below which asserts that (C_4, P_6) -free atoms have bounded clique-width.

► **Theorem 4.** Every (C_4, P_6) -free atom has bounded clique-width.

The proof of Theorem 4 is deferred to subsection 3.3.

Proof of Theorem 3 (assuming Theorem 4). Let G be a (C_4, P_6) -free graph. We find the clique decomposition of Tarjan [49] in $O(mn)$ time and this gives a binary decomposition tree T where the root of T is G and the leaves are induced subgraphs of G without clique cutsets. Tarjan [49] showed that there are at most $O(n)$ leaves and that the chromatic number of any node in T is the maximum of the chromatic numbers of its children. Therefore, determining $\chi(G)$ reduces to determining the chromatic number of atoms. Now it follows from Theorem 4 that each atom has bounded clique-width and thus the chromatic number can be found in polynomial time by Theorem 2. ◀

The remainder of the section is organized as follows. In subsection 3.1, we present the key tools on clique-width that play an important role in the proof of Theorem 4. In subsection 3.2, we list structural properties around a 5-cycle in a (C_4, P_6) -free graph that are frequently used in later proofs. We then present our main proof, the proof of Theorem 4, in subsection 3.3.

3.1 Clique-width

Let $G = (V, E)$ be a graph and H be a proper homogeneous set in G . Then $V \setminus H$ is partitioned into two subsets N and M where N is complete to H and M is anti-complete to H . Let $h \in H$ be an arbitrary vertex and $G_h = G - (H \setminus \{h\})$. We say that H and G_h are *factors* of G with respect to H . Suppose that τ is a k_1 -expression for G_h using labels $1, \dots, k_1$ and σ is a k_2 -expression for H using labels $1, \dots, k_2$. Then substituting $i(h)$ in τ with $\rho_{1 \rightarrow i} \dots \rho_{k_2 \rightarrow i} \sigma$ results in a k -expression for G where $k = \max\{k_1, k_2\}$.

► **Lemma 5** ([12]). The clique-width of any graph G is the maximum clique-width of any prime induced subgraph of G .

A bipartite graph is a *chain* graph if it is $2P_2$ -free. A *co-bipartite chain* graph is the complement of a bipartite chain graph. Let G be a (not necessarily bipartite) graph such that $V(G)$ is partitioned into two subsets A and B . We say that a k -expression for G is *nice* if all vertices in A end up with the same label i and all vertices in B end up with the same label j with $i \neq j$. It is well-known that any co-bipartite chain graph whose vertex set is partitioned into two cliques has a nice 4-expression.

► **Lemma 6** (Folklore). There is a nice 4-expression for any co-bipartite chain graph.

We now use divide-and-conquer to show that a special graph class has clique-width at most 4. This plays a crucial role in our proof of the main theorem (Theorem 4).

► **Lemma 7.** *Let G be a C_4 -free graph such that $V(G)$ is partitioned into two subsets A and B that satisfy the following conditions:*

- (i) A is a clique;
- (ii) B is P_4 -free;
- (iii) no vertex in A has two non-adjacent neighbours in B ;
- (iv) there is no induced P_4 in G that starts with a vertex in A followed by three vertices in B .

Then there is a nice 4-expression for G .

Proof. We use induction on $|B|$. If B contains at most one vertex, then G is a co-bipartite chain graph and the lemma follows from Theorem 6. So, we assume that B contains at least two vertices. Since B is P_4 -free, either B or \bar{B} is disconnected [48]. Suppose first that B is disconnected. Then B can be partitioned into two nonempty subsets B_1 and B_2 that are anti-complete to each other. Let $A_1 = N(B_1) \cap A$ and $A_2 = A \setminus A_1$. Clearly, $G[A_i \cup B_i]$ with the partition (A_i, B_i) satisfies all the conditions of the lemma for each $1 \leq i \leq 2$. Note also that, by (iii), A_1 is anti-complete to B_2 and A_2 is anti-complete to B_1 . By the inductive hypothesis there is a nice 4-expression τ_i for $G[A_i \cup B_i]$ in which all vertices in A_i and B_i have labels 2 and 4, respectively. Now $\rho_{1 \rightarrow 2}(\eta_{1,2}(\tau_1 \oplus \rho_{2 \rightarrow 1}\tau_2))$ is a nice 4-expression for G .


Suppose now that \bar{B} is disconnected. This means that B can be partitioned into two subsets B_1 and B_2 that are complete to each other. Since G is C_4 -free, either B_1 or B_2 is a clique. Without loss of generality, we may assume that B_1 is a clique. Moreover, we choose the partition (B_1, B_2) such that B_1 is maximal. Then every vertex in B_2 is not adjacent to some vertex in B_2 for otherwise we could have moved such a vertex to B_1 . If $B_2 = \emptyset$ then G is a co-bipartite chain graph and so the lemma follows from Theorem 6. Therefore, we assume in the following that $B_1, B_2 \neq \emptyset$. Let $A_1 = N(B_1) \cap A$ and $A_2 = A \setminus A_1$. Note that A_2 is anti-complete to B_1 .

We claim that $N(B_2) \cap A$ is complete to B_1 . Suppose, by contradiction, that $a \in N(B_2) \cap A$ and $b_1 \in B_1$ are not adjacent. By definition, a has a neighbour $b \in B_2$. Recall that b is not adjacent to some vertex $b' \in B_2$. Now a, b, b_1, b' induces either a P_4 or a C_4 , depending on whether a and b' are adjacent. This contradicts (iv) or the C_4 -freeness of G . This proves the claim. Therefore, A_2 is anti-complete to B_2 and $N(B_2) \cap A = N(B_2) \cap A_1$ (see Figure 1). Consequently, $G[(A_1 \cap N(B_2)) \cup B_2]$ with the partition $(A_1 \cap N(B_2), B_2)$ satisfies all the conditions of the lemma. By the inductive hypothesis there is a nice 4-expression τ for $G[(A_1 \cap N(B_2)) \cup B_2]$ in which all vertices in $A \cap N(B_2) = A_1 \cap N(B_2)$ and B_2 have labels 2 and 4, respectively. In addition, note that $(A_1 \setminus N(B_2), B_1)$ is a co-bipartite chain graph. It then follows from Theorem 6 that there is a nice 4-expression ϵ for it in which all vertices in $A_1 \setminus N(B_2)$ and B_1 have labels 1 and 3, respectively. Then

$$\sigma = \rho_{3 \rightarrow 4}(\rho_{1 \rightarrow 2}(\eta_{3,4}(\eta_{2,3}(\eta_{1,2}(\epsilon \oplus \tau))))))$$

is a nice 4-expression for $G - A_2$. Let δ be a 2-expression for A_2 in which all vertices in A_2 have label 1. Then $\rho_{1 \rightarrow 2}(\eta_{1,2}(\delta \oplus \sigma))$ is a nice 4-expression for G . This completes the proof. ◀

Let $G = (V, E)$ be a graph and X and Y two disjoint subsets of $V(G)$. We say that (X, Y) is a *homogeneous pair of sets* in G if no vertex in $V \setminus (X \cup Y)$ distinguishes two vertices in X or in Y . If both X and Y are cliques then (X, Y) is a *homogeneous pair of*



■ **Figure 1** The case \overline{B} is disconnected. Shaded circles represent cliques. A thick line between two sets represents that the two sets are complete, and a dotted line represents that the edges between the two sets can be arbitrary. Two sets are anti-complete if there is no line between them.

cliques. Note that homogeneous sets are special cases of homogeneous pair of sets where one of X and Y is empty. We establish a novel framework via existing results on clique-width for bounding the clique-width of a graph by the clique-width of its subgraphs induced by homogeneous pairs of sets.

► **Lemma 8.** *Let G be a graph such that $V(G)$ can be partitioned into a subset V_0 of vertices of constant size, a constant number of pairs of sets (S_i, T_i) for $1 \leq i \leq r$ and a subset V' of vertices such that*

- (i) *for each $1 \leq i \leq r$, (S_i, T_i) is a homogeneous pair of sets in $G - (V_0 \cup \bigcup_{j=1}^{i-1} (S_j \cup T_j))$;*
- (ii) *for each $1 \leq i \leq r$, $G[S_i \cup T_i]$ has bounded clique-width; and*
- (iii) *$G[V']$ has bounded clique-width.*

Then G has bounded clique-width.


Proof. Let $G_1 = G - V_0$ and $G_{i+1} = G_i - (S_i \cup T_i)$ for $1 \leq i \leq r$. Note that $G_{r+1} = G[V']$. First of all, it follows from [39] that G has bounded clique-width if and only if G_1 has bounded clique-width. In addition, (i) says that (S_i, T_i) is a homogeneous pair of sets in G_i . Let N_i and M_i be sets of vertices in G_i that are complete to S_i and T_i , respectively. For each i we do in G_i two *bipartite complementations* on the pairs $(S_i, V(G_i) \setminus N_i)$ and $(T_i, V(G_i) \setminus M_i)$, which means that we interchange edges and non-edges between the pairs. This results in a graph G' on the same vertex set as G_1 that is the disjoint union of $G[S_i \cup T_i]$ and $G[V']$. It follows from [32] that G_1 has bounded clique-width if and only if each $G[S_i \cup T_i]$ and $G[V']$ have bounded clique-width. Now the lemma follows from our assumptions (ii) and (iii). ◀

3.2 Structure around a 5-Cycle

Let $G = (V, E)$ be a graph and H be an induced subgraph of G . We partition $V \setminus V(H)$ into subsets with respect to H as follows: for any $X \subseteq V(H)$, we denote by $S(X)$ the set of vertices in $V \setminus V(H)$ that have X as their neighbourhood among $V(H)$, i.e.,

$$S(X) = \{v \in V \setminus V(H) : N_{V(H)}(v) = X\}.$$

For $0 \leq j \leq |V(H)|$, we denote by S_j the set of vertices in $V \setminus V(H)$ that have exactly j neighbours among $V(H)$. Note that $S_j = \bigcup_{X \subseteq V(H): |X|=j} S(X)$. We say that a vertex in S_j is a *j -vertex*. Let G be a (C_4, P_6) -free graph and $C = 1, 2, 3, 4, 5$ be an induced C_5 in G . We partition $V \setminus C$ with respect to C as above. All indices below are modulo 5. Since G is C_4 -free, there is no vertex in $V \setminus C$ that is adjacent to vertices i and $i + 2$ but not to vertex $i + 1$. In particular, $S(1, 3)$, S_4 , etc. are empty. The following properties of $S(X)$ were proved in [25] using the fact that G is (C_4, P_6) -free.



■ **Figure 2** The Petersen graph.


- (P1) $S_5 \cup S(i-1, i, i+1)$ is a clique.
- (P2) $S(i)$ is complete to $S(i+2)$ and anti-complete to $S(i+1)$. Moreover, if neither $S(i)$ nor $S(i+2)$ are empty then both sets are cliques.
- (P3) $S(i, i+1)$ is complete to $S(i+1, i+2)$ and anti-complete to $S(i+2, i+3)$. Moreover, if neither $S(i, i+1)$ nor $S(i+1, i+2)$ are empty then both sets are cliques.
- (P4) $S(i-1, i, i+1)$ is anti-complete to $S(i+1, i+2, i+3)$.
- (P5) $S(i)$ is anti-complete to $S(j, j+1)$ if $j \neq i+2$. Moreover, if a vertex in $S(i+2, i+3)$ is not anti-complete to $S(i)$ then it is universal in $S(i+2, i+3)$.
- (P6) $S(i)$ is anti-complete to $S(i+1, i+2, i+3)$.
- (P7) $S(i-2, i+2)$ is anti-complete to $S(i-1, i, i+1)$.
- (P8) Either $S(i)$ or $S(i+1, i+2)$ is empty. By symmetry, either $S(i)$ or $S(i-1, i-2)$ is empty.
- (P9) At least one of $S(i-1, i)$, $S(i, i+1)$ and $S(i+2, i-2)$ is empty.

3.3 Proof of Theorem 4

In this section, we give a proof of Theorem 4. A graph is *chordal* if it does not contain any induced cycle of length at least 4. The following structure of (C_4, P_6) -free graphs discovered by Brandstädt and Hoàng [4] is of particular importance in our proofs below.

- **Theorem 9** ([4]). *Let G be a (C_4, P_6) -free atom. Then the following statements hold:*
 - (i) *every induced C_5 is dominating;*
 - (ii) *if G contains an induced C_6 which is not dominating, then G is the join of a blow-up of the Petersen graph (Figure 2) and a (possibly empty) clique.*

We say that an atom is *strong* if it has no pair of twin vertices or universal vertices. Note that a pair of twin vertices and a universal vertex in a graph give rise to two special kinds of proper homogeneous sets such that one of the factors decomposed by these homogeneous sets is a clique. Therefore, removing twin vertices and universal vertices does not change the clique-width of the graph by Theorem 5. So, to prove Theorem 4 it suffices to prove the theorem for strong atoms. We follow the approach in [18]. In [18], the first and second authors showed how to derive a useful decomposition theorem for (C_4, P_6) -free atoms by eliminating a sequence F_1, C_6, F_2 and C_5 (see Figure 3 for the graphs F_1 and F_2) of induced subgraphs and then employing Dirac's classical theorem [15] on chordal graphs. Here we adopt the same strategy and show in Theorem 10–Theorem 13 below that if a (C_4, P_6) -free strong atom G contains an induced C_5 or C_6 , then it has bounded clique-width. The remaining case is therefore that G is chordal and so G is a clique by Dirac's theorem [15]. Since cliques have clique-width 2, Theorem 4 follows. It turns out that we can easily prove Theorem 10 and Theorem 11 via the framework formulated in Theorem 8 using the structure of the graphs discovered in [18]. The difficulty is, however, that we have to extend the structural analysis



■ **Figure 3** Two special graphs F_1 and F_2 .

in [18] extensively for Theorem 12 and Theorem 13 and provide new insights on bounding the clique-width of certain special graphs using divide-and-conquer (see Theorem 7).

► **Lemma 10.** *If a (C_4, P_6) -free strong atom G contains an induced F_1 , then G has bounded clique-width.*

► **Lemma 11.** *If a (C_4, F_1, P_6) -free strong atom G contains an induced C_6 , then G has bounded clique-width.*

► **Lemma 12.** *If a (C_4, C_6, F_1, P_6) -free strong atom G contains an induced F_2 , then G has bounded clique-width.*

► **Lemma 13.** *If a $(C_4, C_6, F_1, F_2, P_6)$ -free strong atom G contains an induced C_5 , then G has bounded clique-width.*

We illustrate our techniques by giving a proof of Theorem 12 below and omit the proofs of the other lemmas.

Proof of Theorem 12. Let G be a (C_4, C_6, F_1, P_6) -free strong atom that contains an induced subgraph H that is isomorphic to F_2 with $V(H) = \{1, 2, 3, 4, 5, t, x, y\}$ such that $1, 2, 3, 4, 5, 1$ induces the *underlying* 5-cycle C , and t is adjacent to $5, 1$ and 2 , x is adjacent to $4, 5$ and y is adjacent to 2 and 3 . Moreover, t is adjacent to both x and y , see Figure 3. We partition $V(G)$ with respect to C . We choose H such that C has $|S_2|$ maximized. Note that $x \in S(4, 5)$, $y \in S(2, 3)$ and $t \in S(5, 1, 2)$.

The overall strategy is to first decompose G into a subset V_0 of constant size, constant number of homogeneous pairs of sets, and a subset V' , and then finish off the proof via Theorem 8 by showing that each homogeneous pair of sets and $G[V']$ have bounded clique-width where Theorem 7 is employed.

We start with the decomposition. Since $S(2, 3)$ and $S(4, 5)$ are not empty, it follows from (P8) that $S_1 = S(2) \cup S(5)$. If both $S(2)$ and $S(5)$ are not empty, say $u \in S(2)$ and $v \in S(5)$, then $u, 2, 3, 4, 5, v$ induces either a P_6 or a C_6 , depending on whether u and v are adjacent. This shows that $S_1 = S(i)$ for some $i \in \{2, 5\}$. Now we argue that $S_2 = S(2, 3) \cup S(4, 5)$. If $S(3, 4)$ contains a vertex z , then z is adjacent to x and y by (P3) but not adjacent to t by (P7). This implies that t, x, z, y induces a C_4 , so, $S(3, 4) = \emptyset$. If $S(1, 2)$ contains a vertex z , then z is adjacent to y by (P3) and so $1, z, y, 3, 4, 5, 1$ induces a C_6 , a contradiction. This shows that $S(1, 2) = \emptyset$. By symmetry, $S(5, 1) = \emptyset$. Therefore, $S_2 = S(2, 3) \cup S(4, 5)$. The following properties among subsets of G were proved in [18].

- (a) Each vertex in $S(5, 1, 2)$ is either complete or anti-complete to S_2 .
- (b) $S(2, 3)$ and $S(4, 5)$ are cliques.

- (c) Each vertex in $S(3, 4, 5) \cup S(4, 5, 1)$ is either complete or anti-complete to $S(4, 5)$. By symmetry, each vertex in $S(1, 2, 3) \cup S(2, 3, 4)$ is either complete or anti-complete to $S(2, 3)$.
- (d) $S(4, 5)$ is anti-complete to $S(2, 3, 4)$. By symmetry, $S(2, 3)$ is anti-complete to $S(3, 4, 5)$.
- (e) $S(1, 2, 3)$ is complete to $S(5, 1, 2)$. By symmetry, $S(5, 1, 2)$ is complete to $S(4, 5, 1)$.
- (f) $S(4, 5)$ is complete to $S(4, 5, 1)$. By symmetry, $S(2, 3)$ is complete to $S(1, 2, 3)$.
- (g) $S(1, 2, 3)$ is complete to $S(2, 3, 4)$. By symmetry, $S(3, 4, 5)$ is complete to $S(4, 5, 1)$.
- (h) S_5 is complete to S_2 .

Recall that $S_1 = S(i)$ for some $i \in \{2, 5\}$. By symmetry, we may assume that $S_1 = S(5)$. Note that $S(5)$ is complete to $S(4, 5, 1)$ by Theorem 9 and anti-complete to $S(1, 2, 3)$ by (P6). It follows from (P1), (P4), (P7), (e), (f) and (g) that $S(i-1, i, i+1) \cup \{i\}$ is a homogeneous clique in G and therefore $S(i-1, i, i+1) = \emptyset$ for $i = 2, 5$. Similarly, $S(4, 5)$ is a homogeneous clique in G by (P7), (a)-(d), (f) and (h) and so $S(4, 5) = \{x\}$. Let $T = \{t \in S(5, 1, 2) : t \text{ is complete to } S_2\}$.

- (1) $S(5)$ is anti-complete to $S(5, 1, 2) \setminus T$.

Let $u \in S(5)$ and $t' \in S(5, 1, 2) \setminus T$. If u and t' are adjacent, then $u, t', 2, 3, 4, x$ induces either a P_6 or a C_6 , depending on whether u and x are adjacent. ■

By (1) and (d), $(S(5, 1, 2) \setminus T) \cup \{1\}$ is a homogeneous set in G and so $S(5, 1, 2) \setminus T = \emptyset$. In other words, $S(5, 1, 2)$ is complete to S_2 . We now partition $S(5)$ into $X = \{v \in S(5) : v \text{ has a neighbour in } S(2, 3)\}$ and $Y = S(5) \setminus X$.

- (2) X is anti-complete to $S(3, 4, 5)$.

Let $v \in X$ and $s \in S(3, 4, 5)$ be adjacent. By the definition of X , v has a neighbour $y' \in S(2, 3)$. By (d), y' is not adjacent to s and so $v, y', 3, s$ induces a C_4 . ■

- (3) X is complete to $S(5, 1, 2)$.

Assume, by contradiction, that $v \in X$ and $t' \in T$ are not adjacent. By the definition of X , v has a neighbour $y' \in S(2, 3)$. Since t' is adjacent to y' , $v, 5, t', y'$ induces a C_4 . ■

- (4) X is anti-complete to Y .

Suppose that $u \in X$ and $v \in Y$ are adjacent. Let $y' \in S(2, 3)$ be a neighbour of u . Note that x is adjacent to neither u nor v by (P5). But now $x, 4, 3, y', u, v$ induces a P_6 . ■

- (5) X is complete to S_5 .

Suppose that $v \in X$ and $u \in S_5$ are not adjacent. Let $y' \in S(2, 3)$ be a neighbour of v . By (h), y' and u are adjacent. Then $u, 5, v, y'$ induces a C_4 . ■

It follows from (P1)-(P7), (a)-(d), (f), (h) and (2)-(5) that $(X, S(2, 3))$ is a homogeneous pair of sets in G .

- (6) For each connected component A of Y , each vertex in $S(5, 1, 2) \cup S(3, 4, 5)$ is either complete or anti-complete to A .

Let A be an arbitrary component of Y . Suppose that $s \in S(5, 1, 2) \cup S(3, 4, 5)$ distinguishes an edge aa' in A , say s is adjacent to a but not adjacent to a' . We may assume by symmetry that $s \in S(5, 1, 2)$. Then $a', a, s, 2, 3, 4$ induces a P_6 , a contradiction. ■

- (7) Each component of Y has a neighbour in both $S(5, 1, 2)$ and $S(3, 4, 5)$.

Suppose that a component A of Y does not have a neighbour in one of $S(5, 1, 2)$ and $S(3, 4, 5)$, say $S(5, 1, 2)$. Then $S_5 \cup S(3, 4, 5) \cup \{5\}$ is a clique cutset of G by (4). ■

(8) *Each component of Y is a clique.*

Let A be an arbitrary component of Y . By (7), A has a neighbour $s \in S(5, 1, 2)$ and $r \in S(3, 4, 5)$. Note that s and r are not adjacent. Moreover, $\{s, r\}$ is complete to A by (6). Now (8) follows from the fact that G is C_4 -free. ■

(9) *Y is complete to S_5 .*

Suppose, by contradiction, that $v \in Y$ and $u \in S_5$ are not adjacent. By (7), v has a neighbour $s \in S(5, 1, 2)$ and $r \in S(3, 4, 5)$. Then v, s, u, r induces a C_4 . ■

It follows from (P1), (h), (5) and (9) that each vertex in S_5 is a universal vertex in G and so $S_5 = \emptyset$. Let $S'(3, 4, 5) = \{s \in S(3, 4, 5) : s \text{ has a neighbour in } Y\}$ and $S''(3, 4, 5) = S(3, 4, 5) \setminus S'(3, 4, 5)$. Note that $S''(3, 4, 5)$ is anti-complete to Y . We now show further properties of Y and $S'(3, 4, 5)$.

(10) *$S'(3, 4, 5)$ is complete to $S(2, 3, 4)$.*

Suppose, by contradiction, that $r' \in S'(3, 4, 5)$ is not adjacent to $s \in S(2, 3, 4)$. By the definition of $S'(3, 4, 5)$, r' has a neighbour $v \in Y$. Then $v, r', s, 2, 1$ induces a P_6 . ■

(11) *Each vertex in $S(5, 1, 2)$ is either complete or anti-complete to Y .*

Let $t' \in S(5, 1, 2)$ be an arbitrary vertex. Suppose that t' has a neighbour $u \in Y$. Let A be the component of Y containing u . Then t' is complete to A by (6). It remains to show that t' is adjacent to each vertex $u' \in Y \setminus A$. By (7), u has a neighbour $s \in S(3, 4, 5)$. Note that $C' = u, t', y, 3, s$ induces a C_5 . Moreover, x and s are not adjacent for otherwise x, s, u, t' induces a C_4 . This implies that x is adjacent only to t' on C' . On the other hand, u' is not adjacent to any of $u, 3$ and y . This implies that u' is adjacent to either s or t' by Theorem 9. If u' is not adjacent to t' , then u' is adjacent to s . This implies that $u', s, 3, y, t', x$ induces a P_6 or C_6 , depending on whether u' and x are adjacent. Therefore, u' is adjacent to t' . Since u' is an arbitrary vertex in $Y \setminus A$, this proves (11). ■

(12) *$S'(3, 4, 5)$ is anti-complete to x .*

Suppose not. Let $s \in S'(3, 4, 5)$ be adjacent to x . By definition, s has a neighbour $y' \in Y$. Note that x and y' are not adjacent by (P5). By (6) and (7), y' has a neighbour $t \in T = S(5, 1, 2)$. So, t is adjacent to x . But now s, y', t, x induces a C_4 . ■

It follows from (P1)-(P7), (d), (2), (4), (10), (11) and (12) that $(Y, S'(3, 4, 5))$ is a homogeneous pair of sets in G . Let $S'(5, 1, 2) = \{s \in S(5, 1, 2) : s \text{ is complete to } Y\}$. Then $S(5, 1, 2) \setminus S'(5, 1, 2)$ is anti-complete to Y by (11). It follows from (3) that both $S'(5, 1, 2)$ and $S(5, 1, 2) \setminus S'(5, 1, 2)$ are homogeneous cliques in G . So, $|S(5, 1, 2)| \leq 2$. Now $V(G)$ is partitioned into a subset $V_0 = C \cup S(5, 1, 2) \cup \{x\}$ of vertices of size at most 8, two homogeneous pairs of sets $(X, S(2, 3))$ and $(Y, S'(3, 4, 5))$, and a subset $V' = S''(3, 4, 5) \cup S(2, 3, 4)$.

We now apply Theorem 8 to finish off the proof by showing that each of $G[X \cup S(2, 3)]$, $G[Y \cup S'(3, 4, 5)]$, and $G[V']$ has bounded clique-width. First of all, $G[V']$ has clique-width 4 by Theorem 6. Secondly, note that no vertex in $S(1, 2)$ can have two non-adjacent neighbours in X since G is C_4 -free. If there is an induced $P_4 = y', x_1, x_2, x_3$ such that $y' \in S(2, 3)$ and $x_i \in X$, then $x_3, x_2, x_1, y', 3, 4$ induces a P_6 in G . Now if $P = x_1, x_2, x_3, x_4$ is an induced P_4 in $G[X]$, any neighbour y_1 of x_1 is not adjacent to x_3 and x_4 . But then $P \cup \{y_1\}$ contains such a labelled P_4 in $G[X \cup S(2, 3)]$. Therefore, $G[X \cup S(2, 3)]$ with the partition $(X, S(2, 3))$ satisfies all the conditions of Theorem 7 and so has clique-width at most 4. Finally, note that each vertex in $S(3, 4, 5)$ can have neighbours in at most one component of Y due to (7), (11) and the fact that G is C_4 -free. It then follows from (6)-(8) that $G[Y \cup S'(3, 4, 5)]$ with

the partition $(Y, S'(3, 4, 5))$ satisfies all the condition in Theorem 7 (where $A = S'(3, 4, 5)$ and $B = Y$) and so has clique-width at most 4. This completes our proof. ◀

We are now ready to prove our main theorem.

Proof of Theorem 4. Let G be a (C_4, P_6) -free atom. Let G' be the graph obtained from G by removing twin vertices and universal vertices. It follows from Theorem 10–Theorem 13 that if G' contains an induced C_5 or C_6 , then G' has bounded clique-width. Therefore, we can assume that G' is also (C_5, C_6) -free and therefore G' is chordal. It then follows from a well-known result of Dirac [15] that G' is a clique whose clique-width is 2. Finally, $cw(G) = cw(G')$ by Theorem 5 and this completes the proof. ◀

4 The Hardness Result

A graph is a *split graph* if its vertex set can be partitioned into two disjoint sets C and I such that C is a clique and I is an independent set. The pair (C, I) is called a *split partition* of G . A split graph is *complete* if it has a *complete* split partition, that is, a partition (C, I) such that C and I are complete to each other. A *list assignment* of a graph $G = (V, E)$ is a function L that prescribes, for each $u \in V$, a finite list $L(u) \subseteq \{1, 2, \dots\}$ of colours for u . The *size* of a list assignment L is the maximum list size $|L(u)|$ over all vertices $u \in V$. A colouring c *respects* L if $c(u) \in L(u)$ for all $u \in V$. The LIST COLOURING problem is to decide whether a given graph G has a colouring c that respects a given list assignment L . We sketch a proof of our hardness result, in which we construct a graph G' that is neither $(sP_2 + P_8)$ -free nor $(sP_2 + P_4 + P_5)$ -free for any $s \geq 0$. Hence, a different construction is needed for tightening our hardness result (if possible).

► **Theorem 14.** COLOURING is NP-complete for $(C_4, 3P_3, P_3 + P_6, 2P_5, P_9)$ -free graphs.

Proof. (Sketch) We reduce from LIST COLOURING on complete split graphs with a list assignment of size at most 3. It is known that LIST COLOURING is NP-complete for this graph class [20].

Let G be a complete split graph with a list assignment L of size at most 3. From (G, L) we construct an instance (G', k) of COLOURING as follows. Let $k \leq 3|V(G)|$ be the size of the union of all lists $L(u)$. Let (C, I) be a complete split partition of $V(G)$. Let G' be the graph of size $O(|V(G)|k)$ obtained from G as follows. Take a clique X on k vertices x_1, \dots, x_k . For each $u \in V(G)$, introduce a clique Y_u of size $k - |L(u)|$ such that every vertex of Y_u is adjacent to u and to every x_i with $i \in L(u)$ (so, each vertex in every Y_u is adjacent to exactly one vertex of $V(G)$, namely vertex u). By construction, G has a colouring that respects L if and only if G' has a k -colouring. Moreover, it can be readily checked that G' is $(C_4, 3P_3, P_3 + P_6, 2P_5, P_9)$ -free. ◀

5 Conclusions

We gave an almost complete dichotomy for COLOURING restricted to (C_4, P_t) -free graphs and leave open only the cases when $7 \leq t \leq 8$. We believe the techniques developed in this paper could be useful for solving open questions regarding COLOURING on other hereditary graph classes. The natural candidate class for a polynomial-time result of COLOURING is the class of (C_4, P_7) -free graphs. However, this may require significant efforts for the following reason. Lozin and Malyshev [38] determined the complexity of COLOURING for \mathcal{H} -free graphs for every finite set of graphs \mathcal{H} consisting only of 4-vertex graphs except when

\mathcal{H} is $\{K_{1,3}, 4P_1\}$, $\{K_{1,3}, 2P_1 + P_2\}$, $\{K_{1,3}, 2P_1 + P_2, 4P_1\}$ or $\{C_4, 4P_1\}$. Solving any of these open cases would be considered as a major advancement in the area. Since $(C_4, 4P_1)$ -free graphs are (C_4, P_7) -free, polynomial-time solvability of COLOURING on (C_4, P_7) -free graphs implies polynomial-time solvability for COLOURING on $(C_4, 4P_1)$ -free graphs. As a first step, we aim to apply the techniques of this paper to $(C_4, 4P_1)$ -free graphs.

We recall that the complexity of COLOURING on (C_s, P_t) -free graphs is known for all $s \geq 5$ and $t \geq 1$ (Theorem 1) and that the complexity of COLOURING on (C_3, P_t) -free graphs is also known due to the results of [5] and [30] except if $7 \leq t \leq 21$. The class of (C_3, P_7) -free graphs is also a natural class to consider. Interestingly, every (C_3, P_7) -free graph is 5-colourable. This follows from a result of Gravier, Hoàng and Maffray [23] who proved that for any two integers $r, t \geq 1$, every (K_r, P_t) -free graph can be coloured with at most $(t-2)^{r-2}$ colours. On the other hand, 3-COLOURING is polynomial-time solvable for P_7 -free graphs [3]. Hence, in order to solve COLOURING for (C_3, P_7) -free graphs we may instead consider 4-COLOURING for (C_3, P_7) -free graphs. This problem seems also highly non-trivial.

References

- 1 Alexandre Blanché, Konrad K. Dabrowski, Matthew Johnson, and Daniël Paulusma. Hereditary graph classes: When the complexities of Colouring and Clique Cover coincide. *CoRR*, abs/1607.06757, 2016.
- 2 John Adrian Bondy and Uppaluri S.R. Murty. *Graph Theory*, volume 244 of *Springer Graduate Texts in Mathematics*. Springer, 2008.
- 3 Flavia Bonomo, Maria Chudnovsky, Peter Maceli, Oliver Schaudt, Maya Stein, and Mingxian Zhong. Three-coloring and list three-coloring of graphs without induced paths on seven vertices. *Combinatorica*, (in press).
- 4 Andreas Brandstädt and Chinh T. Hoàng. On clique separators, nearly chordal graphs, and the maximum weight stable set problem. *Theoretical Computer Science*, 389:295–306, 2007.
- 5 Andreas Brandstädt, Tilo Klemmt, and Suhail Mahfud. P_6 - and triangle-free graphs revisited: structure and bounded clique-width. *Discrete Mathematics and Theoretical Computer Science*, 8:173–188, 2006.
- 6 Andreas Brandstädt, Van Bang Le, and Jeremy P. Spinrad. *Graph Classes: A Survey*, volume 3 of *SIAM Monographs on Discrete Mathematics and Applications*. Philadelphia, 1999.
- 7 Hajo Broersma, Petr A. Golovach, Daniël Paulusma, and Jian Song. Determining the chromatic number of triangle-free $2P_3$ -free graphs in polynomial time. *Theoretical Computer Science*, 423:1–10, 2012.
- 8 Kathie Cameron and Chinh T. Hoàng. Solving the clique cover problem on (bull, C_4) -free graphs. *CoRR*, abs/1704.00316, 2017.
- 9 Maria Chudnovsky, Peter Maceli, Juraj Stacho, and Mingxian Zhong. 4-Coloring p_6 -free graphs with no induced 5-cycles. *Journal of Graph Theory*, 84:262–285, 2017.
- 10 Maria Chudnovsky and Juraj Stacho. 3-colorable subclasses of P_8 -free graphs. *Manuscript*, 2017.
- 11 Bruno Courcelle, Johann A. Makowsky, and Udi Rotics. Linear time solvable optimization problems on graphs of bounded clique-width. *Theory of Computing Systems*, 33:125–150, 2000.
- 12 Bruno Courcelle and Stephan Olariu. Upper bounds to the clique width of graphs. *Discrete Applied Mathematics*, 101:77–114, 2000.
- 13 Konrad K. Dabrowski, François Dross, and Daniël Paulusma. Colouring diamond-free graphs. *Journal of Computer and System Sciences*, 89:410–431, 2017.

- 14 Konrad K. Dabrowski, Petr A. Golovach, and Daniël Paulusma. Colouring of graphs with Ramsey-type forbidden subgraphs. *Theoretical Computer Science*, 522:34–43, 2014.
- 15 Gabriel A. Dirac. On rigid circuit graphs. *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg*, 25:71–76, 1961.
- 16 Thomas Emden-Weinert, Stefan Hougardy, and Bernd Kreuter. Uniquely colourable graphs and the hardness of colouring graphs of large girth. *Combinatorics, Probability and Computing*, 7:375–386, 1998.
- 17 Wolfgang Espelage, Frank Gurski, and Egon Wanke. How to solve NP-hard graph problems on clique-width bounded graphs in polynomial time. *Proc. WG 2001, LNCS*, 2204:117–128, 2001.
- 18 Serge Gaspers and Shenwei Huang. Linearly χ -bounding (P_6, C_4) -free graphs. *Proc. WG 2017, LNCS*, 10520:263–274, 2017.
- 19 Petr A. Golovach, Matthew Johnson, Daniël Paulusma, and Jian Song. A survey on the computational complexity of coloring graphs with forbidden subgraphs. *Journal of Graph Theory*, 84:331–363, 2017.
- 20 Petr A. Golovach and Daniël Paulusma. List coloring in the absence of two subgraphs. *Discrete Applied Mathematics*, 166:123–130, 2014.
- 21 Petr A. Golovach, Daniël Paulusma, and Jian Song. Coloring graphs without short cycles and long induced paths. *Discrete Applied Mathematics*, 167:107–120, 2014.
- 22 Martin C. Golumbic. *Algorithmic Graph Theory and Perfect Graphs*, volume 57 of *Annals of Discrete Mathematics*. North-Holland Publishing Co., 2004.
- 23 Sylvain Gravier, Chinh T. Hoàng, and Frédéric Maffray. Coloring the hypergraph of maximal cliques of a graph with no long path. *Discrete Mathematics*, 272:285–290, 2003.
- 24 Martin Grötschel, László Lovász, and Alexander Schrijver. Polynomial algorithms for perfect graphs. *Annals of Discrete Mathematics*, 21:325–356, 1984.
- 25 Pavol Hell and Shenwei Huang. Complexity of coloring graphs without paths and cycles. *Discrete Applied Mathematics*, 216, Part 1:211–232, 2017.
- 26 Chinh T. Hoàng, Marcin Kamiński, Vadim V. Lozin, Joe Sawada, and Xiao Shu. Deciding k -colorability of P_5 -free graphs in polynomial time. *Algorithmica*, 57:74–81, 2010.
- 27 Chinh T. Hoàng and D. Adam Lazzarato. Polynomial-time algorithms for minimum weighted colorings of $(P_5, \overline{P_5})$ -free graphs and similar graph classes. *Discrete Applied Mathematics*, 186:106–111, 2015.
- 28 Ian Holyer. The NP-Completeness of edge-coloring. *SIAM Journal on Computing*, 10:718–720, 1981.
- 29 Shenwei Huang. Improved complexity results on k -coloring P_t -free graphs. *European Journal of Combinatorics*, 51:336–346, 2016.
- 30 Shenwei Huang, Matthew Johnson, and Daniël Paulusma. Narrowing the complexity gap for colouring (C_s, P_t) -free graphs. *The Computer Journal*, 58:3074–3088, 2015.
- 31 Tommy R. Jensen and Bjarne Toft. *Graph Coloring Problems*. Wiley Interscience, 1995.
- 32 Marcin Kamiński, Vadim V. Lozin, and Martin Milanič. Recent developments on graphs of bounded clique-width. *Discrete Applied Mathematics*, 157:2747–2761, 2009.
- 33 Thiyagarajan Karthick, Frédéric Maffray, and Lucas Pastor. Polynomial cases for the vertex coloring problem. *CoRR*, abs/1709.07712, 2017.
- 34 Daniel Kobler and Udi Rotics. Edge dominating set and colorings on graphs with fixed clique-width. *Discrete Applied Mathematics*, 126:197–221, 2003.
- 35 Daniel Král', Jan Kratochvíl, Zsolt Tuza, and Gerhard J. Woeginger. Complexity of coloring graphs without forbidden induced subgraphs. *Proc. WG 2001, LNCS*, 2204:254–262, 2001.
- 36 Daniel Leven and Zvi Galil. NP completeness of finding the chromatic index of regular graphs. *Journal of Algorithms*, 4:35–44, 1983.

- 37 László Lovász. Coverings and coloring of hypergraphs. *Congressus Numerantium*, VIII:3–12, 1973.
- 38 Vadim V. Lozin and Dmitriy S. Malyshev. Vertex coloring of graphs with few obstructions. *Discrete Applied Mathematics*, 216, Part 1:273–280, 2017.
- 39 Vadim V. Lozin and Dieter Rautenbach. On the band-, tree-, and clique-width of graphs with bounded vertex degree. *SIAM Journal on Discrete Mathematics*, 18:195–206, 2004.
- 40 Johann A. Makowsky and Udi Rotics. On the clique-width of graphs with few P_4 's. *International Journal of Foundations of Computer Science*, 10:329–348, 1999.
- 41 Dmitriy S. Malyshev. The coloring problem for classes with two small obstructions. *Optimization Letters*, 8:2261–2270, 2014.
- 42 Dmitriy S. Malyshev. Two cases of polynomial-time solvability for the coloring problem. *Journal of Combinatorial Optimization*, 31:833–845, 2016.
- 43 Dmitriy S. Malyshev and O. O. Lobanova. Two complexity results for the vertex coloring problem. *Discrete Applied Mathematics*, 219:158–166, 2017.
- 44 Michael Molloy and Bruce Reed. *Graph Colouring and the Probabilistic Method*. Springer-Verlag Berlin Heidelberg, 2002.
- 45 Sang-Il Oum and Paul D. Seymour. Approximating clique-width and branch-width. *Journal of Combinatorial Theory, Series B*, 96:514–528, 2006.
- 46 Michaël Rao. MSOL partitioning problems on graphs of bounded treewidth and clique-width. *Theoretical Computer Science*, 377:260–267, 2007.
- 47 David Schindl. Some new hereditary classes where graph coloring remains NP-hard. *Discrete Mathematics*, 295:197–202, 2005.
- 48 D. Seinsche. On a property of n -colorable graphs. *Journal of Combinatorial Theory, Series B*, 16:191–193, 1974.
- 49 Robert E. Tarjan. Decomposition by clique separators. *Discrete Mathematics*, 55:221–232, 1985.