

Greedy Bipartite Matching in Random Type Poisson Arrival Model

Allan Borodin

University of Toronto, 10 Kings College Road, Toronto, Canada
bor@cs.toronto.edu

Christodoulos Karavasilis

University of Toronto, 10 Kings College Road, Toronto, Canada
ckar@cs.toronto.edu

Denis Pankratov

University of Toronto, 10 Kings College Road, Toronto, Canada
denisp@cs.toronto.edu

Abstract

We introduce a new random input model for bipartite matching which we call the Random Type Poisson Arrival Model. Just like in the known i.i.d. model (introduced by Feldman et al. [6]), online nodes have types in our model. In contrast to the adversarial types studied in the known i.i.d. model, following the random graphs studied in Mastin and Jaillet [1], in our model each type graph is generated randomly by including each offline node in the neighborhood of an online node with probability c/n independently. In our model, nodes of the same type appear consecutively in the input and the number of times each type node appears is distributed according to the Poisson distribution with parameter 1. We analyze the performance of the simple greedy algorithm under this input model. The performance is controlled by the parameter c and we are able to exactly characterize the competitive ratio for the regimes $c = o(1)$ and $c = \omega(1)$. We also provide a precise bound on the expected size of the matching in the remaining regime of constant c . We compare our results to the previous work of Mastin and Jaillet who analyzed the simple greedy algorithm in the $G_{n,n,p}$ model where each online node type occurs exactly once. We essentially show that the approach of Mastin and Jaillet can be extended to work for the Random Type Poisson Arrival Model, although several nontrivial technical challenges need to be overcome. Intuitively, one can view the Random Type Poisson Arrival Model as the $G_{n,n,p}$ model with less randomness; that is, instead of each online node having a new type, each online node has a chance of repeating the previous type.

2012 ACM Subject Classification Theory of computation → Graph algorithms analysis, Theory of computation → Online algorithms

Keywords and phrases bipartite matching, stochastic input models, online algorithms, greedy algorithms

Digital Object Identifier 10.4230/LIPIcs.APPROX-RANDOM.2018.5

Related Version <https://arxiv.org/abs/1805.00578>

Funding Research is supported by NSERC.

Acknowledgements We thank the anonymous reviewers of APPROX/RANDOM 2018, whose valuable comments helped us improve the presentation of the paper.



© Allan Borodin, Christodoulos Karavasilis, and Denis Pankratov;
licensed under Creative Commons License CC-BY
Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques
(APPROX/RANDOM 2018).

Editors: Eric Blais, Klaus Jansen, José D. P. Rolim, and David Steurer; Article No. 5; pp. 5:1–5:15



Leibniz International Proceedings in Informatics
Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

1 Introduction

Online bipartite matching is a problem with a wide variety of applications and has received significant attention after the seminal work of Karp, Vazirani and Vazirani [10] who showed an optimal $(1 - 1/e)$ randomized algorithm for the unweighted case in the adversarial input model. Applications such as internet advertising (see the excellent survey by Mehta [13] and references therein) and job allocation have given rise to problems that are naturally described as bipartite matching problems and this has been a strong motivation for developing better algorithms. Most of the work in online bipartite matching is with respect to a vertex arrival model where vertices on one side of the bipartite graph are known to the algorithm in advance and vertices on the other side are revealed online along with their adjacent vertices.

Adversarial models are, of course, pessimistic in terms of performance results. For many applications, more realistic assumptions will lead to significantly improved results. In this regard, various stochastic input models have been proposed and analyzed. These include the random order model, known and unknown i.i.d. distribution models, and the Erdős-Rényi random graph model. We propose a new model (see Section 2) that is closely related to both the i.i.d. model and the Erdős-Rényi model. The motivation for our model is that in various applications (e.g. document streams [11] and internet traffic [15]), one often observes the bursty nature of inputs. One can also think of this as a very restricted form of a Markov chain model.

Whereas previous work in the i.i.d. models take advantage of the input distributions in terms of the decisions being made (i.e. as to how to match the online vertices), we follow the work of Mastin and Jaillet [1] who study the performance of the simplest greedy algorithm (which we will call GREEDY) that always matches (when possible) an online vertex to the first (in some fixed ordering of the offline vertices) unmatched adjacent neighbor¹. We note that this simple greedy algorithm does not utilize any information about the nature of the input sequence (including the number of online vertices).

We know that theoretically (i.e. in terms of the expected approximation ratio), that knowledge of the distribution and/or randomization in the algorithm will allow for online algorithms with improved performance. However, in simulations with respect to various stochastic models, we find that the simplest deterministic greedy algorithm is competitive with more specialized algorithms [5]. We also note that when a type graph is chosen adversarially in the i.i.d. model, Goel and Mehta [7] show that the approximation ratio of GREEDY is precisely $1 - 1/e \approx .632$.

Mastin and Jaillet show that the approximation ratio of GREEDY in the Erdős-Rényi model is at least 0.837. Intuitively, our model introduces correlations between consecutive online nodes. This causes significant technical difficulties in the analysis that we are able to overcome. In addition, the correlations do not seem favourable to the GREEDY algorithm, and it is natural to expect GREEDY to have worse performance in our model than in the Erdős-Rényi model. Our results confirm this intuition. Our input model has a parameter c that controls the expected degree of online nodes. Similar to the work of Mastin and Jaillet [1], we analyze the performance of GREEDY in three different regimes of c : $c = o(1)$, $c = \omega(1)$, and constant c . We compute the exact asymptotic fraction of offline nodes matched by GREEDY in each of these regimes. We also show how to derive an upper bound on the

¹ In the version of GREEDY that is studied by Mastin and Jaillet, an online vertex is matched to a randomly chosen available neighbor. Later, we shall see that for our input model, as well as $G_{n,n,p}$, the two versions of GREEDY have the same approximation ratio, since they result in the same stochastic process governing the change in the number of matched offline nodes when an online node arrives.

size of a maximum matching in our model from the existing upper bounds on the size of a maximum matching in the Erdős-Rényi model. Combining the two results, we derive a lower bound on the approximation ratio of GREEDY in all regimes of c . Minimizing this lower bound over c , we find that GREEDY has approximation ratio ≥ 0.715 for all values of c in our model.

Organization. The rest of the paper is organized as follows. Section 2 describes our model and different views of it, as well as some background material needed for the rest of the paper. Section 3 is the technical core of the paper and contains the analysis of GREEDY in the three regimes of c : $c = o(1)$, $c = \omega(1)$, and $c = \Theta(1)$. Section 4 concludes the paper with a brief discussion of various input models and some open questions.

2 Preliminaries

We shall consider bipartite graphs $G = (U, V, E)$ in the vertex arrival model. The vertices in U are referred to as the offline vertices and are known to an algorithm in advance. The vertices in V are referred to as online vertices and arrive one at a time. When a vertex from V is revealed, all its neighbors in U are revealed as well. A simple greedy algorithm matches the arriving vertex $v \in V$ to the first available neighbor in U (if there is one, according to some fixed ordering). We shall denote this algorithm by GREEDY. We consider the behavior of GREEDY on specific families of random graphs generated by, what we call, the Random Type Poisson Arrival Model. This model has two parameters: $n \in \mathbb{N}$ which is equal to $|U|$ and intuitively measures the size of the instance, and $c \in \mathbb{R}_{\geq 0}$ which controls the density of edges. The random graph in our model is generated as follows: the offline nodes U are set to be $[n]$. Online nodes are generated iteratively and randomly using the following process. For each $i \in [n]$ generate a random type i by including each offline node $j \in [n]$ in the neighborhood of the type with probability c/n independently. Then we sample Z_i from the Poisson distribution with parameter 1 and generate Z_i online nodes of type i , and continue. We shall denote a graph G distributed according to the Random Type Poisson Arrival Model with parameters n and c as $G \sim \text{RTPAM}(n, c)$. Our model can be viewed in different ways. We refer to the current view of the $\text{RTPAM}(n, c)$ model as the “one-step view.” Next we describe another view.

The bipartite Erdős-Rényi model is denoted by $G_{n,n,p}$. The random graph $G = (U, V, E)$ in the $G_{n,n,p}$ model is generated as follows: $U = V = [n]$, and each edge $\{i, j\}$ is included in G with probability p independently. Our $\text{RTPAM}(n, c)$ model can be alternatively viewed as a two-step process. At first, we generate a *type graph* $\hat{G} = (U, V, E)$ from the $G_{n,n,p}$ distribution where $p = c/n$. For each type $i \in V$ we sample $Z_i \sim \text{Poi}(1)$. Secondly, an *instance graph* is created by keeping the same U as in the type graph, and replicating each type i node Z_i times (note that this means removing type i when $Z_i = 0$). In the rest of the paper, we shall freely switch between the two different views of the $\text{RTPAM}(n, c)$ model. Thus, we shall occasionally refer to the type graph of the $\text{RTPAM}(n, c)$ graph. We refer to this alternative view of the $\text{RTPAM}(n, c)$ model as the “two-step view.”

Intuitively, in the one-step view, Z_i is drawn from the Poisson distribution in an online fashion whereas in the two-step view, Z_i is drawn initially. It should be clear that these views do not change the model. However, our proofs are facilitated by having these different views.

We shall measure the performance of an algorithm in two ways: in terms of the asymptotic approximation ratio, and in terms of the fraction of the matched offline nodes.

5:4 Greedy Bipartite Matching in RTPAM

► **Definition 1.** Let ALG be a deterministic online algorithm solving the bipartite matching problem over random graphs G_n parameterized by the input size n . We write $\text{ALG}(G_n)$ to denote the size of the matching (random variable) that is constructed by running ALG on G_n . We write $\text{OPT}(G_n)$ to denote the size of a maximum matching in G_n . The *asymptotic approximation ratio* of ALG with respect to G_n is defined as:

$$\rho(\text{ALG}, G_n) = \liminf_{n \rightarrow \infty} \frac{\mathbb{E}(\text{ALG}(G_n))}{\mathbb{E}(\text{OPT}(G_n))}.$$

The *fraction of matched offline nodes* of ALG with respect to G_n is defined as:

$$\mu(\text{ALG}, G_n) = \liminf_{n \rightarrow \infty} \frac{\mathbb{E}(\text{ALG}(G_n))}{n}.$$

We shall use the following notation for the well-known distributions:

► **Definition 2.**

- $\text{Poi}(\lambda)$ – the Poisson distribution with parameter λ ,
- $\text{Bin}(n, p)$ – the Binomial distribution with parameters $n \in \mathbb{Z}_{\geq 0}$ (number of trials) and $p \in [0, 1]$ (probability of success)
- $\text{Ber}(p)$ – the Bernoulli distribution with parameter p .

We shall also write $\text{Poi}(\lambda)$, $\text{Bin}(n, p)$, etc., as a placeholder for a random variable distributed according to the corresponding distribution. This is done to simplify the notation when the name of the random variable is not important and only the parameters of the distribution are of interest.

► **Definition 3.** We write $\mathbb{I}(E)$ to denote the indicator random variable for an event E .

In the paper, we show how to compute $\mu(\text{GREEDY}, \text{RTPAM}(n, c))$ exactly. In order to provide a bound on the asymptotic approximation ratio of GREEDY we also need to know the size of a maximum matching in $\text{RTPAM}(n, c)$. The size of a maximum matching is not known even in $G_{n,n,c/n}$ model, but there is a known upper bound due to Bollobás and Brightwell that we will be able to use to derive a lower bound on the asymptotic approximation ratio of GREEDY in the $\text{RTPAM}(n, c)$ model.

► **Theorem 4 ([4]).**

$$\frac{\mathbb{E}(\text{OPT}(G_{n,n,c/n}))}{n} \leq 2 - \frac{\gamma^* + \gamma_* + \gamma^* \gamma_*}{c} + o(1),$$

where γ_* is the smallest solution to the equation $x = c \exp(-c \exp(-x))$ and $\gamma^* = c \exp(-\gamma_*)$. See Figure 4 for the shape of this upper bound as a function of c .

We shall later compare our results with the following result giving the performance of GREEDY in the $G_{n,n,c/n}$ model due to Mastin and Jaillet.

► **Theorem 5 ([1]).** For $c \in \mathbb{R}_{>0}$ we have

$$\mu(\text{GREEDY}, G_{n,n,c/n}) = 1 - \frac{\log(2 - e^{-c})}{c}.$$

3 Greedy Matching in Random Type Poisson Arrival Model

3.1 The Regime of $c = o(1)$

In this section we show that in expectation GREEDY finds an almost-maximum matching in RTPAM(n, c) when $c = o(1)$. The high level idea is to consider the two-step view of RTPAM(n, c) and observe that when $c = o(1)$ most of the type graph consists of *isolated* edges – edges with both endpoints of degree 1. Thus, no matter how many times an online node corresponding to an isolated edge is generated in the instance graph, both GREEDY and OPT can match it exactly once (given that it is generated at all). The expected number of the non-isolated edges is of smaller order of magnitude than the expected number of isolated edges and can be ignored for the purpose of computing the asymptotic approximation ratio.

► **Theorem 6.** *Let $c = o(1)$ then we have:*

$$\rho(\text{GREEDY}, \text{RTPAM}(n, c)) = 1.$$

Proof. (similar to the proof of Lemma 2 in [1]) Set $p = c/n = o(1/n)$, and consider the two-step view of RTPAM(n, c). The probability that an edge between i and j appears and is isolated in *the type graph* is:

$$\Pr(\{i, j\} \text{ is isolated in the type graph}) = p(1-p)^{2n-2}.$$

Observe that if type j is generated at least once in *the instance graph*, i.e., $Z_j \geq 1$, and $\{i, j\}$ is an isolated edge in *the type graph* then GREEDY will include $\{i, j\}$ in the matching exactly once. In addition, observe that the events $Z_j \geq 1$ and “ $\{i, j\}$ is isolated in the type graph” are independent. Let $W_{i,j}$ be a random variable indicating whether $\{i, j\}$ is included by GREEDY. Then, we have:

$$\Pr(W_{i,j} = 1) \geq \Pr(Z_j \geq 1) \Pr(\{i, j\} \text{ is isolated in the type graph}) = \left(1 - \frac{1}{e}\right) p(1-p)^{2n-2}.$$

Let M be the matching produced by GREEDY. We have $|M| = \sum_{i,j} W_{i,j}$. It follows that:

$$\mathbb{E}(|M|) = \mathbb{E}\left(\sum_{i,j} W_{i,j}\right) = \sum_{i,j} \Pr(W_{i,j} = 1) \geq n^2 \left(1 - \frac{1}{e}\right) p(1-p)^{2n-2}.$$

Let Q_j denote the number of neighbors of a node of type j . Observe that in the instance graph the nodes corresponding to type j can be matched in an optimal matching at most $Q_j \mathbb{I}(Z_j \geq 1)$ times. Note that Q_j and Z_j are independent. Let M^* denote a maximum matching in the instance graph. Then we have

$$\mathbb{E}(|M^*|) \leq \sum_j \mathbb{E}(Q_j \mathbb{I}(Z_j \geq 1)) = \sum_j np(1 - 1/e) = n^2 p(1 - 1/e).$$

Combining this with the above lower bound on $\mathbb{E}(|M|)$ we get

$$\frac{\mathbb{E}(|M|)}{\mathbb{E}(|M^*|)} \geq (1-p)^{2n-2} = \left(e^{-o(\frac{1}{n})}\right)^{2n-2} = e^{-o(1)} \xrightarrow{n \rightarrow \infty} 1. \quad \blacktriangleleft$$

3.2 The Regime of Constant c

Fix a constant $c \in \mathbb{R}_{>0}$. We can view GREEDY as constructing the matching in rounds. Consider the one-step view of RTPAM(n, c). During round i , a new type is created and Z_i nodes corresponding to that type are generated. We let Y_i denote the number of online nodes matched by GREEDY by the beginning of round i . We also let X_i denote the number of neighbors of type i that were *not* matched in any of the earlier rounds. In this section, we show how to compute the asymptotic fraction of matched offline nodes by GREEDY *exactly*. More specifically, we derive an asymptotically accurate (implicit) expression for Y_n and show how to compute it for each value of c . In addition, we show that existing upper bounds on the maximum matching in the $G_{n,n,c/n}$ model carry over to the RTPAM(n, c) model. This allows us to derive lower bounds on the *competitive ratio* of GREEDY in RTPAM(n, c).

High level idea. We use the method of partial differential equations (see, e.g., [19, 18, 12]) to derive the asymptotic behavior of Y_n . The goal is to write the expression $\mathbb{E}(Y_{i+1} - Y_i \mid Y_i)$ in terms of Y_i/n , i.e., $\mathbb{E}(Y_{i+1} - Y_i \mid Y_i) = f_c(Y_i/n)$ for some “simple” function f_c . This gives us a difference equation for Y_i . Now, pretend that there is a function $g_c : [0, 1] \rightarrow [0, 1]$ that gives a good approximation to Y_i , i.e., $g_c(t) \approx Y_{tn}/n$ for $t \in [0, 1]$. Consider a syntactic replacement of the difference equation for Y_i with a differential equation for g_c , i.e., $g'_c(t) = f_c(g_c(t))$. In addition, set the correct initial value condition $g_c(0) = Y_0 = 0$. The differential equation method allows us to conclude that under a mild condition on f_c (namely, being Lipschitz), the solution g_c is unique and asymptotically converges to Y_n/n , i.e., $Y_n = g_c(1)n + o(n)$. In particular, we have $\mu(\text{GREEDY}, \text{RTPAM}(n, c)) = g_c(1)$. In our setting, we will see that it is not clear how to write $\mathbb{E}(Y_{i+1} - Y_i \mid Y_i)$ as a function of Y_i/n . It turns out that the method still works as long as $\mathbb{E}(Y_{i+1} - Y_i \mid Y_i)$ is close to $f_c(Y_i/n)$ in the following sense: $\lim_{n \rightarrow \infty} |\mathbb{E}(Y_{i+1} - Y_i \mid Y_i) - f_c(Y_i/n)| = 0$. This is precisely what we do in this section.

The number of nodes matched by GREEDY in round i is exactly equal² to $\min(X_i, Z_i)$. By the definition of the RTPAM(n, c) model, we have $X_i \sim \text{Bin}(n - Y_i, c/n)$. Therefore, we have

$$\mathbb{E}(Y_{i+1} - Y_i \mid Y_i) = \mathbb{E}(\min(\text{Bin}(n - Y_i, c/n), \text{Poi}(1)) \mid Y_i). \quad (1)$$

Unfortunately, as mentioned above this expectation does not seem to have a nice form and we do not know how to set up an associated differential equation. Instead, we shall approximate the difference equation $\mathbb{E}(Y_{i+1} - Y_i \mid Y_i)$ by another expression that is easier to handle. Those familiar with the Poisson limit theorem will immediately recognize the following as the most natural choice:

$$\mathbb{E}(\min(\text{Bin}(n - Y_i, c/n), \text{Poi}(1)) \mid Y_i) \approx \mathbb{E}(\min(\text{Poi}(c(1 - Y_i/n)), \text{Poi}(1)) \mid Y_i).$$

We need to analyze how accurate this approximation is, but first we show how to derive a relatively simple expression for the right hand side. Define $h(x) := \mathbb{E}(\min(\text{Poi}(x), \text{Poi}(1)))$. Although the function $h(x)$ does not have a closed-form expression in terms of widely-known functions such as \sin, \cos, \exp , etc., it does have a closed-form expression in terms of the modified Bessel functions of the first kind and the Marcum’s Q functions:

² Observe that this holds in all regimes of c and that this expression remains the same if we consider the version of GREEDY of Mastin and Jaillet, where an online node is matched to a *random* available neighbor. Therefore, the exact version of GREEDY does not matter for our input model. Similar argument applies to $G_{n,n,p}$.

► **Definition 7.** The modified Bessel functions of the first kind are defined as follows:

$$I_k(x) = \sum_{i=0}^{\infty} \frac{1}{i! \Gamma(i+k+1)} \left(\frac{x}{2}\right)^{2i+k}.$$

For an integer $k \geq 0$, it becomes $I_k(x) = \sum_{i=0}^{\infty} \frac{1}{i!(i+k)!} \left(\frac{x}{2}\right)^{2i+k}$. We note that the modified Bessel functions have the following symmetry property: $I_k(x) = I_{-k}(x)$.

► **Definition 8.** Marcum’s Q function is defined as follows:

$$Q_n(a, b) = \exp\left(-\frac{a^2 + b^2}{2}\right) \sum_{k=1-n}^{\infty} \left(\frac{a}{b}\right)^k I_k(ab).$$

Now, the closed-form expression for $h(x)$ can be derived from the answer on Stats Stackexchange [16]. For completeness, we reproduce the derivation in Appendix A.

► **Lemma 9** ([16]). *For $x > 0$ we have*

$$h(x) = \frac{1 + x - 2e^{-x-1} (I_0(2\sqrt{x}) + \sqrt{x}I_1(2\sqrt{x})) - (1-x)(1 - 2Q_1(\sqrt{2x}, \sqrt{2}))}{2}.$$

Thus, in terms of our overview we have $f_c(x) = h(c(1-x))$, because we hope to show that

$$\mathbb{E}(\min(\text{Bin}(n-Y_i, c/n), \text{Poi}(1)) \mid Y_i) \approx \mathbb{E}(\min(\text{Poi}(c(1-Y_i/n)), \text{Poi}(1)) \mid Y_i) = h(c(1-Y_i/n)).$$

To apply the method of differential equations we need to analyze the above approximation and show that f_c is Lipschitz. We start by analyzing how good the approximation is.

► **Lemma 10.**

$$\lim_{n \rightarrow \infty} \left| \mathbb{E}(\min(\text{Bin}(n - Y_i, c/n), \text{Poi}(1)) \mid Y_i) - h(c(1 - Y_i/n)) \right| = 0.$$

Proof. We introduce the following useful notation: $N = n - Y_i$, $p = c/n$, and $\lambda = c(1 - Y_i/n)$. Moreover, we let $b_r = \Pr(\text{Bin}(N, p) = r)$ and $p_r(a) = \Pr(\text{Poi}(a) = r)$. Let $W_i \sim \text{Poi}(c(1 - Y_i/n))$. Then the statement of the lemma can be translated into

$$\left| \mathbb{E}(\min(X_i, Z_i) - \min(W_i, Z_i) \mid Y_i) \right|$$

The expectation is just a big sum. Let’s consider individual terms and their contributions to the overall sum.

1. **Case: $W_i < X_i < Z_i$.** The contribution of $X_i = k, W_i = j, Z_i = \ell$ when $j < k < \ell$ is $k - j$.
2. **Case: $X_i < W_i < Z_i$.** The contribution of $X_i = j, W_i = k, Z_i = \ell$ when $j < k < \ell$ is $j - k = -(k - j)$.
3. **Case: $W_i < Z_i \leq X_i$.** The contribution of $X_i = k, W_i = j, Z_i = \ell$ when $j < \ell$ and $k \geq \ell$ is $\ell - j$.
4. **Case: $X_i < Z_i \leq W_i$.** The contribution of $X_i = j, W_i = k, Z_i = \ell$ when $j < \ell$ and $k \geq \ell$ is $j - \ell$.
5. **Cases: (a) $X_i = W_i$; (b) $Z_i \leq X_i, W_i$.** The contribution is 0.

5:8 Greedy Bipartite Matching in RTPAM

We pair up terms corresponding to (1) with (2) and terms corresponding to (3) with (4). Define

$$S_1 = \sum_{k,j,\ell=0}^{\infty} \mathbb{I}(j < k < \ell) ((k-j)b_k p_j(\lambda) p_\ell(1) + (j-k)b_j p_k(\lambda) p_\ell(1)),$$

and

$$S_2 = \sum_{k,j,\ell=0}^{\infty} \mathbb{I}(j < \ell, k \geq \ell) ((\ell-j)b_k p_j(\lambda) p_\ell(1) + (j-\ell)b_j p_k(\lambda) p_\ell(1)).$$

Thus, we get that

$$|\mathbb{E}(\min(X_i, Z_i) - \min(W_i, Z_i) \mid Y_i)| = |S_1 + S_2| \leq |S_1| + |S_2|.$$

We will show that $\lim_{n \rightarrow \infty} |S_1| = 0$. Similar argument implies that $\lim_{n \rightarrow \infty} |S_2| = 0$.

We have

$$\begin{aligned} S_1 &= \sum_{k,j,\ell=0}^{\infty} \mathbb{I}(j < k < \ell) p_\ell(1) (k-j)(b_k p_j(\lambda) - b_j p_k(\lambda)) \\ &= \sum_{k,j,\ell=0}^{\infty} \mathbb{I}(j < k < \ell) p_\ell(1) (k-j) ((b_k - p_k(\lambda)) p_j(\lambda) - (b_j - p_j(\lambda)) p_k(\lambda)) \\ &= S_1^1 + S_1^2, \end{aligned}$$

where $S_1^1 = \sum_{k,j,\ell=0}^{\infty} \mathbb{I}(j < k < \ell) p_\ell(1) (k-j)(b_k - p_k(\lambda)) p_j(\lambda)$ and $S_1^2 = S_1 - S_1^1$. Again, $|S_1| \leq |S_1^1| + |S_1^2|$. We will show that $\lim_{n \rightarrow \infty} |S_1^1| = 0$, and a similar argument implies that the same holds for $|S_1^2|$. We have

$$|S_1^1| \leq \sum_{k=0}^{\infty} \left(\sum_{j,\ell=0}^{\infty} \mathbb{I}(j < k < \ell) (k-j) p_\ell(1) p_j(\lambda) \right) |b_k - p_k(\lambda)|.$$

By [14], we have $\lim_{n \rightarrow \infty} |S_1^1| = 0$ if and only if

$$\sum_{k,j,\ell=0}^{\infty} \mathbb{I}(j < k < \ell) (k-j) p_\ell(1) p_j(\lambda) p_k(\lambda) = O(1).$$

Lastly, we have

$$\begin{aligned} \sum_{k,j,\ell=0}^{\infty} \mathbb{I}(j < k < \ell) (k-j) p_\ell(1) p_j(\lambda) p_k(\lambda) &\leq \sum_{k,j,\ell=0}^{\infty} \mathbb{I}(j < k < \ell) k p_\ell(1) p_j(\lambda) p_k(\lambda) \\ &\leq \sum_{k,j,\ell=0}^{\infty} p_\ell(1) p_j(\lambda) k p_k(\lambda) \leq \lambda \xrightarrow{n \rightarrow \infty} c = O(1). \end{aligned} \quad \blacktriangleleft$$

Due to space considerations we prove that f_c is Lipschitz in Appendix A.

► **Lemma 11.** *The function $f_c(x)$ is Lipschitz on $[0, 1]$.*

Finally, we have all the necessary ingredients to prove the main theorem of this section. Although this theorem does not give an explicit closed-form expression for $\mu(\text{GREEDY, RTPAM}(n, c))$, it gives a simple way to evaluate it numerically for any value of $c > 0$.

► **Theorem 12.**

$$\mu(\text{GREEDY}, \text{RTPAM}(n, c)) = g_c(1),$$

where g_c is a solution to the following differential equation:

$$\begin{aligned} g'_c(t) &= h(c(1 - g_c(t))), \\ g_c(0) &= 0, \end{aligned}$$

and h is given in Lemma 9.

Proof. This is a direct application of Wormald's theorem (Theorem 5.1 in [19]) using Lemmas 10 and 11. ◀

Next, we show two upper bounds on $\mathbb{E}(\text{OPT}(\text{RTPAM}(n, c)))$. The minimum of the two will be used to compute the approximation ratio of GREEDY.

► **Theorem 13.** For all $c \in \mathbb{R}_{>0}$ we have

$$\mathbb{E}(\text{OPT}(\text{RTPAM}(n, c))) \leq \min \left(nc \left(1 - \frac{1}{e} \right), \left(2 - \frac{\gamma^* + \gamma_* + \gamma^* \gamma_*}{c} \right) n + o(n) \right),$$

where γ_* is the smallest solution to the equation $x = c \exp(-c \exp(-x))$ and $\gamma^* = c \exp(-\gamma_*)$.

Proof. The first argument in the minimum follows from the proof of Theorem 6. Let Q_j denote the number of neighbors of a node of type j . Observe that the number of nodes of type j participating in any matching is bounded above by $Q_j \mathbb{I}(Z_j \geq 1)$. Using the fact that Q_j and $\mathbb{I}(Z_j \geq 1)$ are independent, taking the expectation and summing over all j results in the upper bound of $nc \left(1 - \frac{1}{e} \right)$.

The second argument in the minimum follows from the observation

$$\mathbb{E}(\text{OPT}(\text{RTPAM}(n, c))) \leq \mathbb{E}(\text{OPT}(G_{n,n,c/n})) \tag{2}$$

and Theorem 4. Let $\alpha(G)$ denote the independence number of the graph. By Gallai's and König's theorems it suffices to prove that

$$\mathbb{E}(\alpha(\text{RTPAM}(n, c))) \geq \mathbb{E}(\alpha(G_{n,n,c/n})).$$

Consider the two-step view of $\text{RTPAM}(n, c)$. In the first step, a type graph $\widehat{G} = (U, V, E)$ is generated from the distribution $G_{n,n,c/n}$. Let S be a largest independent set in the type graph. Write $S = S_1 \cup S_2$, where $S_1 = S \cap U$ consists of offline nodes and $S_2 = S \cap V$ consists of online types. In the *instance* graph all nodes from S_1 together with all nodes with types from S_2 will form an independent set. In other words, even if a node of a given type is repeated multiple times from S_2 , it can be safely included in an independent set. Thus, we have

$$\begin{aligned} \mathbb{E}(\alpha(\text{RTPAM}(n, c)) \mid S_1, S_2) &\geq \mathbb{E} \left(|S_1| + \sum_{j \in S_2} Z_j \mid S_1, S_2 \right) \\ &= |S_1| + \sum_{j \in S_2} \mathbb{E}(Z_j \mid S_1, S_2) = |S_1| + |S_2| = |S|, \end{aligned}$$

where the last equality follows because Z_j is independent of S_1, S_2 . Taking the expectation over S proves $\mathbb{E}(\alpha(\text{RTPAM}(n, c))) \geq \mathbb{E}(\alpha(G_{n,n,c/n}))$, since $|S|$ has the same distribution as $\alpha(G_{n,n,c/n})$. ◀

3.3 The Regime of $c = \omega(1)$

In this section we show that GREEDY matches almost all offline nodes in RTPAM(n, c) model when $c = \omega(1)$. Consider the two-step view of RTPAM(n, c). Recall that Z_j refers to the number of nodes of type j that are generated, and that X_j refers to the number of neighbors of a node of type j that have not been matched in any of the previous rounds. Also, recall that GREEDY matches $\mathbb{E}(\min(X_j, Z_j))$ in round j in expectation. We will show that in most rounds $\min(X_j, Z_j)$ is very close to Z_j . This finishes the argument, since $\sum_j \mathbb{E}(Z_j) = n$ is the total number of offline nodes and a trivial upper bound on the size of a maximum matching.

► **Theorem 14.** *Let $c = \omega(1)$ then we have:*

$$\rho(\text{GREEDY}, \text{RTPAM}(n, c)) = 1.$$

Proof. Let $p = \frac{c}{n}$ and $k = \frac{n}{\sqrt{c}}$. Fix a round i and assume that at least k offline nodes have not been matched in earlier rounds. Then variable X_i has binomial distribution with at least k trials and the probability of success c/n . We will consider $\tilde{X}_i \sim \text{Bin}(k, c/n)$ such that $\tilde{X}_i \leq X_i$. We will show that $\Pr(\tilde{X}_i \geq Z_i) = 1 - o(1)$. Since $Z_i \sim \text{Poi}(1)$ we have:

$$\Pr(Z_i \geq c^{1/100}) = \frac{1}{e} \sum_{j=c^{1/100}}^{\infty} \frac{1}{j!} \leq \frac{1}{c^{1/100}!} \leq \frac{1}{2c^{1/100}}.$$

For \tilde{X}_i we have $\text{Var}(\tilde{X}_i) = kp(1-p) = \sqrt{c}(1-c/n)$ and $\mathbb{E}(\tilde{X}_i) = \sqrt{c}$. By Chebyshev's inequality

$$\Pr(|\tilde{X}_i - \sqrt{c}| \geq c^{1/3}) \leq \frac{\sqrt{c}(1-c/n)}{c^{2/3}} = \frac{1-c/n}{c^{1/6}}.$$

From these two bounds, it follows that

$$\Pr(\tilde{X}_i \geq Z_i) \geq \Pr(Y_i \leq c^{1/100} \wedge \tilde{X}_i \geq \sqrt{c} - c^{1/3}) \geq 1 - \frac{1-c/n}{c^{1/6}} - \frac{1}{2c^{1/100}} = 1 - o(1).$$

In addition, it is easy to see that in the first $n - 10k$ rounds GREEDY matches at most $n - k$ offline nodes with probability $1 - o(1)$. In particular, the probability of matching more than that is bounded by the probability that $\sum_{i=1}^{n-10k} Z_i > n - k$. Thus, we can condition on having at least k available offline nodes during each of the first $n - 10k$ rounds. Therefore, the expected size of the matching constructed by GREEDY is at least $\sum_{i=1}^{n-10k} \mathbb{E}(\min(Z_i, X_i)) \geq (n - 10k)(1 - o(1)) = n - o(n)$. ◀

3.4 Putting it together

In this section, we take a closer look at our results for GREEDY in RTPAM(n, c) model. We already know that GREEDY achieves competitive ratio 1 in the regimes $c = o(1)$ and $c = \omega(1)$. Hence, we concentrate on the regime of constant c . In Figure 1 we plot the asymptotic fraction of matched offline nodes by GREEDY (Theorem 12) and the upper bound on the fraction of offline nodes in a maximum matching (Theorem 13) as functions of c .

By taking the ratio of the two curves in Figure 1 we obtain a lower bound on the asymptotic approximation ratio of GREEDY in the RTPAM(n, c) model as a function of c . We plot this lower bound in Figure 2. We see that the lower bound achieves a unique minimum on the interval $(0, \infty)$ and that it converges to 1 as c goes to infinity. By numerically minimizing the lower bound we obtain that the minimum of this curve is achieved at $c \approx 0.667766$ and the lower bound is ≈ 0.715071 . Thus, we have the following corollary:

► **Corollary 15.** *For all regimes of c we have*

$$\rho(\text{GREEDY}, \text{RTPAM}(n, c)) \geq 0.715.$$

The shape of the lower bound graph in Figure 2 is a bit strange and it suggests that our lower bound might not be tight. Therefore, we conjecture that it should be possible to strengthen the upper bound on the size of a maximum matching in $\text{RTPAM}(n, c)$.

It is also interesting to compare the performance of GREEDY on $\text{RTPAM}(n, c)$ inputs with its performance on $G_{n,n,c/n}$ inputs. As stated in the introduction, we expect GREEDY to perform worse in the $\text{RTPAM}(n, c)$ model, because the $\text{RTPAM}(n, c)$ model has “less randomness” in the sense of introducing correlations between consecutive online nodes that are not present in the $G_{n,n,c/n}$ model. Indeed, this intuition turns out to be correct. We plot the performance of GREEDY in two models in Figure 3 and observe that GREEDY on $G_{n,n,c/n}$ inputs outperforms $\text{RTPAM}(n, c)$ for constant c .

4 Conclusion

We have introduced a new stochastic model for the online bipartite matching problem. In our model, a random Erdős-Rényi type graph is generated first. Then an input instance graph is generated in rounds where in the i^{th} round, the corresponding input type node appears consecutively n_i times where n_i is distributed according to the Poisson distribution with parameter 1.

More generally, this model is just a specific case of a broad class of stochastic online models for graph problems where type graphs are generated by some random or adversarial processes and then the i^{th} input type node occurs consecutively n_i times where n_i is determined by another random process so as to model some limited form of “locality of reference”. More generally, we could use a Markov process to generate the next type node instance.

The $G_{n,n,p}$ Erdős-Rényi graphs (where $n_i = 1$ for all i) and i.i.d. models where the type graph G is determined adversarially or according to a random process (and where the i^{th} round is drawn i.i.d. from the type graph) fit within this general class of stochastic models.

As in Mastin and Jaillet [1] and Besser and Poloczek [3], we analyze the performance of the simplest greedy algorithm. As in other such studies, it is often the case that simple greedy or “greedy-like” algorithms perform well on real benchmarks or stochastic settings, well beyond what worst case analysis might suggest. Our specific RTPAM model introduces dependencies between online nodes that do not appear in other stochastic models for maximum bipartite matching. These dependencies in the RTPAM model result in some technical challenges in addition to the non-trivial analysis in Mastin and Jaillet. As in Mastin and Jaillet, our analysis falls into three classes depending on the edge probabilities $p = c/n$. As in Mastin and Jaillet, the regimes $c = o(1)$ and $c = \omega(1)$ result in approximation ratios that approach 1 as n increases. And as in Mastin and Jaillet, we obtain an almost precise approximation ratio (modulo the estimate of the expected size of optimal matching) for the regime of constant c . Given the input dependencies our worst case bound (i.e. for the c^* that minimizes the approximation ratio) is significantly less than in Mastin and Jaillet.

As we have suggested, our RTPAM model is just a specific case of a wide class of online stochastic models that have not been studied with respect to any algorithm. We believe that such a study will be both technically interesting as well as becoming more applicable to many “real-world” settings where there is “locality of reference”. Finally, we have begun an experimental study of the performance of Greedy in comparison to algorithms that exploit the underlying type graph in a distributional model (e.g., [6, 17, 2, 9, 5]).

References

- 1 P. Jaillet A. Mastin. Greedy online bipartite matching on random graphs, 2013. URL: <https://arxiv.org/abs/1307.2536v1>.
- 2 M. Zadimoghaddam B. Haeupler, V.S. Mirrokni. Online stochastic weighted matching: Improved approximation algorithms, WINE 2011.
- 3 Bert Besser and Matthias Poloczek. Greedy matching: Guarantees and limitations. *Algorithmica*, 77(1):201–234, 2017.
- 4 Bela Bollobás and Graham Brightwell. The width of random graph orders. *The Mathematical Scientist*, 20, 01 1995.
- 5 Allan Borodin, Christodoulos Karavasilis, and Denis Pankratov. An experimental study of algorithms for online bipartite matching, Unpublished work in progress, 2018.
- 6 Jon Feldman, Aranyak Mehta, Vahab Mirrokni, and S. Muthukrishnan. Online stochastic matching: Beating $1-1/e$. In *Proc. of FOCS*, pages 117–126, 2009.
- 7 Gagan Goel and Aranyak Mehta. Online budgeted matching in random input models with applications to adwords. In *Proc. of SODA*, pages 982–991, 2008.
- 8 Carl W. Helstrom. *Statistical Theory of Signal Detection (Second Edition, Revised and Enlarged)*. International Series of Monographs in Electronics and Instrumentation. Pergamon, second edition, revised and enlarged edition, 1968.
- 9 Patrick Jaillet and Xin Lu. Online stochastic matching: New algorithms with better bounds. *Mathematics of Operations Research*, 39(3):624–646, 2014.
- 10 R. M. Karp, U. V. Vazirani, and V. V. Vazirani. An optimal algorithm for on-line bipartite matching. In *Proc. of STOC*, pages 352–358, 1990.
- 11 Jon M. Kleinberg. Bursty and hierarchical structure in streams. *Data Min. Knowl. Discov.*, 7(4):373–397, 2003.
- 12 Thomas G. Kurtz. Solutions of ordinary differential equations as limits of pure jump markov processes. *Journal of Applied Probability*, 7(1):49–58, 1970.
- 13 A. Mehta. Online matching and ad allocation, *Theoretical Computer Science*, 8(4):265–368, 2012.
- 14 Gordon Simons and N. L. Johnson. On the convergence of binomial to poisson distributions. *Ann. Math. Statist.*, 42(5):1735–1736, 10 1971. doi:10.1214/aoms/1177693172.
- 15 Kannikar Siritwong, Lester Lipsky, and Reda A. Ammar. Study of bursty internet traffic. In *Sixth IEEE International Symposium on Network Computing and Applications (NCA 2007), 12 - 14 July 2007, Cambridge, MA, USA*, pages 53–60, 2007.
- 16 Stats Stackexchange. What is the expectation of the absolute value of the skellam distribution? <https://stats.stackexchange.com/questions/279220/what-is-the-expectation-of-the-absolute-value-of-the-skellam-distribution>. Accessed: 2018-04-11.
- 17 A. Saberi V. H. Manshadi, S. Oveis Gharan. Online stochastic matching: Online actions based on offline statistics, *Mathematics of Operations Research*, 37(4):559-573, 2012.
- 18 Nicholas C. Wormald. Differential equations for random processes and random graphs. *Ann. Appl. Probab.*, 5(4):1217–1235, 11 1995.
- 19 Nicholas C. Wormald. The differential equation method for random graph processes and greedy algorithms. In *Lectures on approximation and randomized algorithms*, pages 73–155, 1999.

A Two Technical Lemmas

In this appendix, we prove two lemmas that are used in Section 3. The first lemma gives a closed-form expression for $h(x) := \mathbb{E}(\min(\text{Poi}(x), \text{Poi}(1)))$ in terms of the modified Bessel functions of the first kind and the Marcum's Q functions. The derivation relies on the answer from Stats Stackexchange [16].

► **Lemma 16** (Lemma 9 restated,[16]). *For $x > 0$ we have*

$$h(x) = \frac{1 + x - 2e^{-x-1} (I_0(2\sqrt{x}) + \sqrt{x}I_1(2\sqrt{x})) - (1-x) (1 - 2Q_1(\sqrt{2x}, \sqrt{2}))}{2}.$$

Proof. A random variable that is equal to the difference between two Poisson random variables has Skellam distribution. As the first step, we reduce the computation of $h(x)$ to the computation of the expectation of an absolute value of a Skellam distributed random variable:

$$\begin{aligned} h(x) &= \mathbb{E}(\min(\text{Poi}(x), \text{Poi}(1))) = \frac{\mathbb{E}(\text{Poi}(x)) + \mathbb{E}(\text{Poi}(1)) - \mathbb{E}(|\text{Poi}(x) - \text{Poi}(1)|)}{2} \\ &= \frac{1 + x - \mathbb{E}(|\text{Poi}(x) - \text{Poi}(1)|)}{2}. \end{aligned}$$

In the rest of the proof, we show how to compute $\mathbb{E}(|\text{Poi}(x) - \text{Poi}(1)|)$. From the PMF of a Skellam variable, one easily obtains the PMF of the absolute value of our Skellam variable:

$$\Pr(|\text{Poi}(x) - \text{Poi}(1)| = k) = \begin{cases} e^{-1-x} (x^{k/2} I_k(2\sqrt{x}) + x^{-k/2} I_{-k}(2\sqrt{x})) & \text{if } k > 0, \\ e^{-1-x} I_0(2\sqrt{x}) & \text{if } k = 0. \end{cases}$$

Then we write down the MGF and simplify it using the Marcum's Q function to get:

$$\begin{aligned} M_{|\text{Poi}(x) - \text{Poi}(1)|}(t) &= e^{-1-x} \left(Q_1(\sqrt{2 \exp(-t)}, \sqrt{2x \exp(t)}) \exp(xe^t + e^{-t}) \right. \\ &\quad \left. + Q_1(\sqrt{2x \exp(-t)}, \sqrt{2 \exp(t)}) \exp(e^t + xe^{-t}) - I_0(2\sqrt{x}) \right). \end{aligned}$$

Now, we can take the derivative of the MGF at $t = 0$ to derive:

$$\mathbb{E}(|\text{Poi}(x) - \text{Poi}(1)|) = 2e^{-x-1} (I_0(2\sqrt{x}) + \sqrt{x}I_1(2\sqrt{x})) + (1-x) (1 - 2Q_1(\sqrt{2x}, \sqrt{2})).$$

◀

In the next lemma we prove that the function defining the differential equation in Section 3 is Lipschitz. This is needed to establish the main result of the paper via Wormald's theorem.

► **Lemma 17** (Lemma 11 restated). *The function $f_c(x)$ is Lipschitz on $[0, 1]$.*

Proof. We prove the statement by showing that the derivative of $f_c(x)$ is bounded on $[0, 1]$. By definition of f_c we have $f_c(x) = h(c(1-x))$. Thus, $f'_c(x) = -ch'(c(1-x))$. Hence bounding $|f'_c(x)|$ on $[0, 1]$ amounts to bounding $h'(x)$ on $[0, c]$. We compute the derivative of h as follows:

$$\begin{aligned} 2h'(x) &= 2 + 2e^{-1-x} (I_0(2\sqrt{x}) + \sqrt{x}I_1(2\sqrt{x})) - e^{-1-x} (3I_1(2\sqrt{x})/\sqrt{x} + I_0(2\sqrt{x}) + I_2(2\sqrt{x})) \\ &\quad - 2Q_1(\sqrt{2x}, \sqrt{2}) + 2(1-x) (Q_2(\sqrt{2x}, \sqrt{2}) - Q_1(\sqrt{2x}, \sqrt{2})) \\ &= 2 + 2e^{-1-x} (I_0(2\sqrt{x}) + \sqrt{x}I_1(2\sqrt{x})) - e^{-1-x} (3I_1(2\sqrt{x})/\sqrt{x} + I_0(2\sqrt{x}) + I_2(2\sqrt{x})) \\ &\quad - 2Q_1(\sqrt{2x}, \sqrt{2}) + 2(1-x) e^{-1-x} I_1(2\sqrt{x})/\sqrt{x} \\ &= e^{-1-x} (I_0(2\sqrt{x}) - I_1(2\sqrt{x})/\sqrt{x} - I_2(2\sqrt{x})) + 2 - 2Q_1(\sqrt{2x}, \sqrt{2}), \end{aligned}$$

where the second equation follows from the definition of the Marcum's Q function and the third equation follows by collecting and simplifying terms with the factor of e^{-1-x} together. We complete the proof of the lemma by bounding each of the terms.


By the definition of the Bessel functions of the first kind we have

$$\begin{aligned} I_0(2\sqrt{x}) - I_1(2\sqrt{x})/\sqrt{x} - I_2(2\sqrt{x}) &= \sum_{i=0}^{\infty} \frac{1}{i!i!} x^i - \sum_{i=0}^{\infty} \frac{1}{i!(i+1)!} x^i - \sum_{i=1}^{\infty} \frac{1}{(i-1)!(i+1)!} x^i \\ &= 1 - 1 + \sum_{i=1}^{\infty} \left(\frac{1}{i!i!} - \frac{1}{i!(i+1)!} - \frac{1}{(i-1)!(i+1)!} \right) x^i \\ &= \sum_{i=1}^{\infty} \frac{1}{i!(i+1)!} x^i \leq \sum_{i=0}^{\infty} \frac{1}{i!} x^i = e^x. \end{aligned}$$


Thus, the first term is bounded by e^c . The second term $2 - 2Q_1(\sqrt{2x}, \sqrt{2})$ is bounded by 2. This follows from interpretation of the Marcum's Q function as a probability – in particular, we have $Q_1(\sqrt{2x}, \sqrt{2}) \in [0, 1]$ (see [8]). All in all, we have $|f'_c(x)| = O(1)$ for $x \in [0, 1]$. ◀

B Figures


In this appendix, we collect all figures mentioned in the paper.




■ **Figure 1** The exact asymptotic ratio of GREEDY to n from Theorem 12, and the upper bound on the asymptotic ratio of a size of maximum matching to n from Theorem 13. Both results are plotted as functions of c in the RTPAM(n, c) model.



■ **Figure 2** The lower bound on the competitive ratio for GREEDY in RTPAM(n, c) model as a function of c .



■ **Figure 3** The exact asymptotic ratio of GREEDY to n in $\text{RTPAM}(n, c)$ input model (Theorem 12) versus the exact asymptotic ratio of GREEDY to n in $G_{n, n, c/n}$ input model (Theorem 5).



■ **Figure 4** The upper bound on the asymptotic ratio of the size of a maximum matching in $G_{n, n, c/n}$ to n given in Theorem 4.