

Improved Strong Spatial Mixing for Colorings on Trees

Charilaos Efthymiou

Department of Computer Science, University of Warwick, UK
charilaos.efthymiou@warwick.ac.uk

Andreas Galanis

Department of Computer Science, University of Oxford, UK
andreas.galanis@cs.ox.ac.uk

Thomas P. Hayes

Department of Computer Science, University of New Mexico, Albuquerque, NM, USA
hayes@cs.unm.edu

Daniel Štefankovič

Department of Computer Science, University of Rochester, NY, USA
stefanko@cs.rochester.edu

Eric Vigoda

School of Computer Science, Georgia Institute of Technology, Atlanta, GA, USA
ericvigoda@gmail.com

Abstract

Strong spatial mixing (SSM) is a form of correlation decay that has played an essential role in the design of approximate counting algorithms for spin systems. A notable example is the algorithm of Weitz (2006) for the hard-core model on weighted independent sets. We study SSM for the q -colorings problem on the infinite $(d+1)$ -regular tree. Weak spatial mixing (WSM) captures whether the influence of the leaves on the root vanishes as the height of the tree grows. Jonasson (2002) established WSM when $q > d + 1$. In contrast, in SSM, we first fix a coloring on a subset of internal vertices, and we again ask if the influence of the leaves on the root is vanishing. It was known that SSM holds on the $(d + 1)$ -regular tree when $q > \alpha d$ where $\alpha \approx 1.763\dots$ is a constant that has arisen in a variety of results concerning random colorings. Here we improve on this bound by showing SSM for $q > 1.59d$. Our proof establishes an L^2 contraction for the BP operator. For the contraction we bound the norm of the BP Jacobian by exploiting combinatorial properties of the coloring of the tree.

2012 ACM Subject Classification Mathematics of computing \rightarrow Discrete mathematics; Theory of computation \rightarrow Random walks and Markov chains

Keywords and phrases colorings, regular tree, spatial mixing, phase transitions

Digital Object Identifier 10.4230/LIPIcs.APPROX-RANDOM.2019.48

Category RANDOM

Funding *Charilaos Efthymiou*: Supported by the Centre of Discrete Mathematics and its Applications (DIMAP), University of Warwick, EPSRC award EP/D063191/1.

Andreas Galanis: The research leading to these results has received funding from the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013) ERC grant agreement no. 334828. The paper reflects only the authors' views and not the views of the ERC or the European Commission. The European Union is not liable for any use that may be made of the information contained therein.

Thomas P. Hayes: Partially supported by NSF CAREER award CCF-1150281.

Daniel Štefankovič: Research supported in part by NSF grant CCF-1563757.

Eric Vigoda: Research supported in part by NSF grants CCF-1617306 and CCF-1563838.



© Charilaos Efthymiou, Andreas Galanis, Thomas P. Hayes, Daniel Štefankovič, and Eric Vigoda; licensed under Creative Commons License CC-BY

Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques (APPROX/RANDOM 2019).

Editors: Dimitris Achlioptas and László A. Végh; Article No. 48; pp. 48:1–48:16



Leibniz International Proceedings in Informatics

Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

1 Introduction

Consider random q -colorings of the complete tree T_h of height h with branching factor d . Does the influence of the leaves on the root decay to zero in the limit as the height grows? If so, this corresponds to weak spatial mixing, which we will define more precisely momentarily.

Now suppose we fix the coloring τ for a subset of internal vertices. Is it still the case that the influence of the leaves on the root decay to zero as the height grows? One might intuitively expect that these internal “agreements” defined by τ only help in the sense that the influence of the leaves decrease, however this problem is much more challenging; it corresponds to strong spatial mixing, which is the focus of this paper.

For statistical physics models, the key algorithmic problems are the counting problem of estimating the partition function and the problem of sampling from the Gibbs distribution, which corresponds to the equilibrium state of the system. Strong spatial mixing (SSM) is a key property of the system for the design of efficient counting/sampling algorithms.

SSM has a variety of algorithmic implications. A direct consequence of SSM on amenable graphs, such as the integer lattice \mathbb{Z}^d , is fast mixing of the Glauber dynamics, which is the simple Markov chain that updates the spin at a randomly chosen vertex in each step, see, e.g. [21, 22, 6, 9, 14, 4, 3]. SSM also plays a critical role in the efficiency of correlation-decay techniques of Weitz [26] which yields an FPTAS for the partition function of the hard-core model in the tree uniqueness region; this approach has been extended to 2-spin antiferromagnetic models [18] and other interesting examples, e.g., [19]; note, the approach of Barvinok [1] utilizing a zero-free region of the partition function in the complex plane has recently been extended to the same range of parameters for the hard-core model [23, 25].

The fundamental question in statistical physics is the uniqueness/non-uniqueness phase transition which corresponds to whether long-range correlations persist or die off, in the limit as the volume of the system tends to infinity. In the uniqueness region the correlations die off, which corresponds to *weak spatial mixing* (WSM). While WSM (or equivalently uniqueness) is a notoriously challenging problem on the 2-dimensional integer lattice \mathbb{Z}^2 (e.g., see the recent breakthrough work of Beffara and Duminil-Copin [2] for the ferromagnetic Potts model), the corresponding WSM problem on the infinite $(d+1)$ -regular tree \mathbb{T}_d , known as the Bethe lattice, is typically simpler since it can be analyzed using recursions due to the absence of cycles (e.g., see Kelly [17] for the hard-core model). However, for the colorings problem, which is the focus of this paper, even WSM is far from trivial on the regular tree [16]. In fact, for the closely related antiferromagnetic Potts model the precise range of parameters for WSM is only known for fixed values of q, d [11].

The focus of this paper is on these correlation decay properties on the infinite $(d+1)$ -regular tree \mathbb{T}_d for the *colorings* problem. We give an informal definition of WSM and SSM, and refer the interested reader to Section 2 for formal definitions.

Let T_h denote the complete tree of height h where all internal vertices have degree $d+1$. For integer $q \geq 3$, let μ_h denote the uniform distribution over proper (vertex) q -colorings of T_h . Consider a pair of sequences of colorings (η_h) and (η'_h) for the leaves of T_h . Let p_h and p'_h denote the marginal probability that the root receives a specific color c under μ_h conditional on the leaves having the fixed coloring η_h and η'_h , respectively. Roughly, if $\lim_{h \rightarrow \infty} |p_h - p'_h| = 0$ for all sequences $(\eta_h), (\eta'_h)$ and colors c , then we say WSM holds (see also Section 2). Jonasson [16] proved that WSM holds when $q \geq d+2$. When $q \leq d+1$, the pair of boundary conditions can actually “freeze” the color at the root; moreover, Brightwell and Winkler [5] showed that there are multiple semi-translation invariant Gibbs measures on \mathbb{T}_d when $q \leq d$.

Now consider an arbitrary coloring τ for a subset $S \subset \mathbb{T}_d$. Let r_h and r'_h denote the marginal probability that the root receives color c under μ_h conditional on $\eta_h \cup \tau$ and $\eta'_h \cup \tau$, respectively. If these limits are the same then we say **SSM** holds. The challenge of establishing **SSM** is illustrated by the fact that if **WSM** holds then we know that $\lim_{h \rightarrow \infty} p_h = 1/q$ but that is not necessarily the case in the **SSM** setting.

Ge and Štefankovič [13] proved that **SSM** holds on \mathbb{T}_d when $q > \alpha d$ where $\alpha \approx 1.763\dots$ is the root of $\frac{1}{\alpha} \exp(1/\alpha) = 1$. Gamarnik, Katz, and Misra [12] extended this result to arbitrary triangle-free graphs of maximum degree d , under the same condition on q . Recent work of Liu, Sinclair, and Srivastava [20] builds upon [12] together with the approximate counting approach of [1, 23] to obtain an **FPTAS** for counting colorings of triangle-free graphs when $q > \alpha d$. Prior to these works, Goldberg, Martin, and Paterson [14] established the above form of **SSM** on triangle-free amenable¹ graphs, also when $q > \alpha d$. In addition to the above results, the threshold $\alpha \approx 1.76\dots$ has arisen in numerous rapid mixing results, e.g., [7, 15, 8].

Our main result presents the first substantial improvement on the 1.76... threshold of [13], we establish **SSM** on the tree when $q > 1.59d$. We state a somewhat informal version of our main theorem here, the formal version will be given once we define more precisely **SSM**, cf. Theorem 3 below.

► **Theorem 1** (Informal version of Theorem 3). *There exists an absolute constant $\beta > 0$ such that, for all positive integers q, d satisfying $q \geq 1.59d + \beta$, the q -coloring model exhibits strong spatial mixing on the regular tree \mathbb{T}_d .*

We remark that the constant 1.59 in Theorem 1 can be replaced with any $\alpha' > 1$ satisfying

$$\frac{1}{\alpha'} \exp\left(\frac{1}{\alpha'}\right) \exp\left(-\frac{1}{\alpha' - 1 + \exp\left(\frac{1}{\alpha' - 1}\right)}\right) < 1,$$

the smallest such value up to four decimal digits is 1.5897.

We give an overview of our proof approach in Section 3 after formally defining **SSM** in Section 2 and stating the formal version of Theorem 1. We then present detailed proofs of the three main lemmas in Section 3.

2 Definitions

Let $q \geq 3$ be an integer and $G = (V, E)$ be a graph. A proper q -coloring of G is an assignment $\sigma : V \rightarrow [q]$ such that for every $(u, v) \in E$ it holds that $\sigma(u) \neq \sigma(v)$. We use Ω_G to denote the set of all proper q -colorings of G and μ_G to denote the uniform probability distribution on Ω_G (provided that Ω_G is non-empty).

For $\sigma \in \Omega_G$ and a set $\Lambda \subset V$, we use σ_Λ to denote the restriction of σ to Λ . When Λ consists of a single vertex v , we will often use the shorthand σ_v to denote the color of v under σ . We say that an assignment $\eta : \Lambda \rightarrow [q]$ is *extendible* if there exists a coloring $\sigma \in \Omega_G$ such that $\sigma_\Lambda = \eta$.

We can now formally define **SSM**.

► **Definition 2.** *Let $\zeta : \mathbb{Z}_{\geq 0} \rightarrow [0, 1]$ be a real-valued function on the positive integers.*

*The q -coloring model exhibits strong spatial mixing, denoted **SSM**, on a finite graph $G = (V, E)$ with decay rate $\zeta(\cdot)$ iff for every $v \in V$, for every $\Lambda \subset V$, for any two extendible*

¹ Roughly, a graph is amenable if for every subset S of vertices, the neighborhood satisfies $|N(S)| \leq \text{poly}(|S|)$.

48:4 Improved Strong Spatial Mixing for Colorings on Trees

assignments $\eta, \eta' : \Lambda \rightarrow [q]$ and any color $c \in [q]$ it holds that

$$\left| \mu_G(\sigma_v = c \mid \sigma_\Lambda = \eta) - \mu_G(\sigma_v = c \mid \sigma_\Lambda = \eta') \right| \leq \zeta(\text{dist}(v, \Delta)), \quad (1)$$

where $\Delta \subseteq \Lambda$ denotes the set of vertices where η and η' disagree.

In the case where G is infinite, we say that the q -coloring model exhibits strong spatial mixing on G with decay rate $\zeta(\cdot)$ if it exhibits strong spatial mixing on every finite subgraph of G with decay rate $\zeta(\cdot)$.

The definition of weak spatial mixing has one modification: in the RHS of (1) we replace $\text{dist}(v, \Delta)$ by the weaker condition $\text{dist}(v, \Lambda)$. WSM says that the influence of a pair of boundary conditions decays at rate $\zeta(\cdot)$ in the distance to the boundary Λ . In SSM the pair of boundaries η, η' might only differ on a subset $\Delta \subset \Lambda$; do these fixed “agreements” on $\Lambda \setminus \Delta$ influence the marginal at v ? If SSM holds then the difference in the marginal at v decays at rate $\zeta(\cdot)$ in the distance to the “disagreements” in η, η' .

With these definitions in place, we are now ready to give the formal version of Theorem 1.

► **Theorem 3.** *There exists an absolute constant $\beta > 0$ such that, for all positive integers q, d satisfying $q \geq 1.59d + \beta$, the q -coloring model exhibits strong spatial mixing on the regular tree \mathbb{T}_d with exponentially decaying rate.*

That is, there exist constants $\alpha, C > 0$ and a function ζ satisfying $\zeta(\ell) \leq C \exp(-\alpha\ell)$ for all integers $\ell \geq 0$ such that for all finite subtrees T of \mathbb{T}_d the q -coloring model exhibits strong spatial mixing on T with decay rate ζ .

3 Proof Approach

For a set $\Lambda \subset V$ and an extendible assignment $\eta : \Lambda \rightarrow [q]$, we use $\pi_{G, v, \eta}$ to denote the q -dimensional probability vector whose entries give the marginal distribution of colors at v under the boundary condition η , i.e., for a color $c \in [q]$, the c -th entry of $\pi_{G, v, \eta}$ is given by $\mu_G(\sigma_v = c \mid \sigma_\Lambda = \eta)$.

The key ingredient to prove Theorem 3 is the following.

► **Theorem 4.** *There exist absolute constants $\beta > 0$ and $U \in (0, 1)$ such that the following holds for all positive integers q, d satisfying $q \geq 1.59d + \beta$.*

Let $T = \mathbb{T}_{d, h, \rho}$ be the d -ary tree with height h rooted at ρ , Λ be a subset of the vertices of T , and $\eta, \eta' : \Lambda \rightarrow [q]$ be two extendible assignments of T with $\text{dist}(\rho, \Delta) \geq 3$ where $\Delta \subseteq \Lambda$ is the set of vertices where η and η' disagree. Let v_1, \dots, v_d be the children of ρ and for $i \in [d]$ let $T_i = (V_i, E_i)$ be the subtree of T rooted at v_i which consists of all descendants of v_i in T . Then

$$\|\pi - \pi'\|_2^2 \leq U \max_{i \in [d]} \|\pi_i - \pi_{i'}\|_2^2,$$

where $\pi = \pi_{T, \rho, \eta}$, $\pi' = \pi_{T, \rho, \eta'}$ and for $i \in [d]$ we denote $\pi_i = \pi_{T_i, v_i, \eta(\Lambda \cap V_i)}$, $\pi_{i'} = \pi_{T_i, v_i, \eta'(\Lambda \cap V_i)}$.

Intuitively, Theorem 4 says that disagreements between η and η' have smaller impact on the marginals as we move upwards on the tree. More precisely, the marginals of the root under η and under η' are closer in L^2 distance than the distance between the marginals of any child (under the induced distributions on the subtrees hanging from them).

Using Theorem 4, the proof of Theorem 3 of strong spatial mixing follows from rather standard considerations, the proof can be found in Section 7. In the following section, we focus on the more interesting proof of Theorem 4 and explain the new aspects of our analysis.

3.1 The three main lemmas

In this section, we lay down the main technical steps in proving Theorem 4. In particular, we will assume throughout that, for appropriate integers q, d, h , $T = \hat{T}_{d,h,\rho}$ is the d -ary tree with height h rooted at ρ , Λ is a subset of the vertices of T , and $\eta, \eta' : \Lambda \rightarrow [q]$ are two extendible assignments of T with $\text{dist}(\rho, \Delta) \geq 3$ where $\Delta \subseteq \Lambda$ is the set of vertices where η and η' disagree. We also let v_1, \dots, v_d be the children of ρ and for $i \in [d]$ let $T_i = (V_i, E_i)$ be the subtree of T rooted at v_i which consists of all descendants of v_i in T .

To prove Theorem 4, we will use tree recursions to express the marginal at the root in terms of the marginals at the children (as in previous works on WSM/SSM, see, e.g., [5, 13, 11]). This recursion is the well-known *Belief Propagation* (BP) equation [24]; our proof of Theorem 4 will be based on bounding appropriately the gradient of the BP equations. The new ingredient in our analysis is that we incorporate the combinatorial structure of agreements close to the root into a refined L^2 analysis of the gradient.

Prior to delving into the analysis, we first describe the BP equation for the colorings model. Following the notation of Theorem 4, let $\boldsymbol{\pi} = \boldsymbol{\pi}_{T,\rho,\eta}$, $\boldsymbol{\pi}' = \boldsymbol{\pi}_{T,\rho,\eta'}$ be the marginal distributions at the root of the tree T under the boundary conditions η and η' , respectively. Similarly, for $i \in [d]$, let $\boldsymbol{\pi}_i, \boldsymbol{\pi}'_i$ be the marginals at the root v_i of the subtree T_i under $\eta(\Lambda \cap V_i)$ and $\eta'(\Lambda \cap V_i)$, respectively.

We can now relate the distribution $\boldsymbol{\pi}$ with the distributions $\{\boldsymbol{\pi}_i\}_{i \in [d]}$ (and similarly, $\boldsymbol{\pi}'$ with the distributions $\{\boldsymbol{\pi}'_i\}_{i \in [d]}$) as follows. For q -dimensional probability vectors $\mathbf{x}_1, \dots, \mathbf{x}_d$ and a color $c \in [q]$, let f_c be the function

$$f_c(\mathbf{x}_1, \dots, \mathbf{x}_d) = \frac{\prod_{i \in [d]} (1 - x_{i,c})}{\sum_{j \in [q]} \prod_{i \in [d]} (1 - x_{i,j})}, \tag{2}$$

where, for $i \in [d]$ and $j \in [q]$, $x_{i,j}$ denotes the j -th entry of the vector \mathbf{x}_i . Then, with π_c and π'_c denoting the c -th entries of $\boldsymbol{\pi}$ and $\boldsymbol{\pi}'$, we have that

$$\begin{aligned} \pi_c &= \mu_T(\sigma_\rho = c \mid \sigma_\Lambda = \eta) = f_c(\boldsymbol{\pi}_1, \dots, \boldsymbol{\pi}_d), \\ \pi'_c &= \mu_T(\sigma_\rho = c \mid \sigma_\Lambda = \eta') = f_c(\boldsymbol{\pi}'_1, \dots, \boldsymbol{\pi}'_d). \end{aligned} \tag{3}$$

The functions $\{f_c\}_{c \in [q]}$ correspond to the BP equations for the coloring model.

We are now ready to describe in more detail our SSM analysis. Specifically, to get a bound on the norm $\|\boldsymbol{\pi} - \boldsymbol{\pi}'\|_2$, we will study the gradient of f_c as we change the arguments $(\boldsymbol{\pi}_1, \dots, \boldsymbol{\pi}_d)$ to $(\boldsymbol{\pi}'_1, \dots, \boldsymbol{\pi}'_d)$ along the line connecting them. Our gradient analysis will take account of the following combinatorial notions.

► **Definition 5.** A vertex v of T is called *frozen under η* if $v \in \Lambda$ and non-frozen otherwise. For a non-frozen vertex v of T , a color k is *blocked for v (under η)* if there is a neighbor $u \in \Lambda$ of v such that $\eta(u) = k$; the color is called *available for v* otherwise.

► **Observation 6.** In the setting of Theorem 4, we have that the disagreements between η and η' occur at distance at least 3 from the root. It follows that the set of the root's children that are frozen as well as the set of blocked colors for each of the non-frozen children are identical under both η and η' .

We will utilize that the gradient components that correspond to either frozen children or blocked colors can be disregarded since, by Observation 6, the corresponding arguments in (3) are fixed to the same value. Namely, we will track, for each color c , the fraction of non-frozen children which have color c available. This will allow us in the upcoming Lemma 10 to aggregate accurately the gradient components corresponding to color c . The following definitions setup some relevant notation.

► **Definition 7.** Let D be the indices of the children of the root which are non-frozen under η and η' . For a color $c \in [q]$, let $\gamma_c \in [0, 1]$ be the fraction of indices $i \in D$ such that color c is available for v_i under η and η' (cf. Observation 6). Let $\boldsymbol{\gamma}$ and $\sqrt{\boldsymbol{\gamma}}$ be the q -dimensional vector with entries $\{\gamma_c\}_{c \in [q]}$ and $\{\sqrt{\gamma_c}\}_{c \in [q]}$, respectively.

Intuitively, if γ_c is close to 0, color c is blocked at a lot of the children and hence the distance $\|\boldsymbol{\pi} - \boldsymbol{\pi}'\|_2$ at the root should not depend a lot on the color c (since most components of the gradient corresponding to color c are zero).

The following couple of definitions will be relevant for capturing more precisely the gradient of the functions $\{f_c\}_{c \in [q]}$. To begin with, the gradient will actually turn out to be related to the value of f_c as we move along the line $(\boldsymbol{\pi}_1, \dots, \boldsymbol{\pi}_d)$ to $(\boldsymbol{\pi}'_1, \dots, \boldsymbol{\pi}'_d)$. More precisely, we have the following definition.

► **Definition 8.** For $t \in [0, 1]$, let $\hat{\boldsymbol{\pi}}(t) = \{\hat{\pi}_c(t)\}_{c \in [q]}$ be the q -dimensional probability vector whose c -th entry is given by $f_c(t\boldsymbol{\pi}_1 + (1-t)\boldsymbol{\pi}'_1, \dots, t\boldsymbol{\pi}_d + (1-t)\boldsymbol{\pi}'_d)$.

Note that $\hat{\boldsymbol{\pi}}(1) = \boldsymbol{\pi}$ and $\hat{\boldsymbol{\pi}}(0) = \boldsymbol{\pi}'$; in this sense, we can think of the vector $\hat{\boldsymbol{\pi}}(t)$ as having the marginals at the root as we interpolate between $(\boldsymbol{\pi}_1, \dots, \boldsymbol{\pi}_d)$ to $(\boldsymbol{\pi}'_1, \dots, \boldsymbol{\pi}'_d)$.

The next definition will be relevant for bounding the L^2 norm of the gradient along the line connecting to $(\boldsymbol{\pi}_1, \dots, \boldsymbol{\pi}_d)$ to $(\boldsymbol{\pi}'_1, \dots, \boldsymbol{\pi}'_d)$. The bound will be in terms of the “marginals” at the root, as captured by the vector $\hat{\boldsymbol{\pi}}(t)$ (cf. Definition 8), and the availability of the q colors at the children, as captured by the vector $\boldsymbol{\gamma}$ (cf. Definition 7). In particular, we will be interested in the L^2 norm of the following matrix, which is an idealized version to the Jacobian of the BP equation (see (13) for the precise formula).²

► **Definition 9.** Let $\hat{\boldsymbol{\pi}}, \hat{\boldsymbol{\gamma}}$ be q -dimensional vectors with non-negative entries. The matrix $\mathbf{M}_{\hat{\boldsymbol{\pi}}, \hat{\boldsymbol{\gamma}}}$ corresponding to the vectors $\hat{\boldsymbol{\pi}}, \hat{\boldsymbol{\gamma}}$ is given by $(\text{diag}(\hat{\boldsymbol{\pi}}) - \hat{\boldsymbol{\pi}}\hat{\boldsymbol{\pi}}^\top)\text{diag}(\hat{\boldsymbol{\gamma}})$.³

Our first main lemma shows how to bound the distance between the marginals at the root under η and η' , i.e., $\|\boldsymbol{\pi} - \boldsymbol{\pi}'\|_2^2$, in terms of the aggregate distance at the children. The new ingredient in our bound is to account more carefully for the availability of the colors at the children (i.e., the vector $\boldsymbol{\gamma}$).

► **Lemma 10.** Let q, d be positive integers so that $q \geq d + 2$. Then

$$\|\boldsymbol{\pi} - \boldsymbol{\pi}'\|_2^2 \leq |D|K^2 \sum_{i \in [d]} \|\boldsymbol{\pi}_i - \boldsymbol{\pi}'_i\|_2^2 \quad \text{where } K := \frac{1}{1 - \frac{1}{q-d}} \max_{t \in (0,1)} \|\mathbf{M}_{\hat{\boldsymbol{\pi}}(t), \sqrt{\boldsymbol{\gamma}}}\|_2,$$

where $D, \boldsymbol{\gamma}, \sqrt{\boldsymbol{\gamma}}$ are as in Definition 7, $\hat{\boldsymbol{\pi}}(t)$ is as in Definition 8, and $\mathbf{M}_{\hat{\boldsymbol{\pi}}(t), \sqrt{\boldsymbol{\gamma}}}$ is as in Definition 9.

Given Lemma 10, we are left with obtaining a good upper bound on the norm $\|\mathbf{M}_{\hat{\boldsymbol{\pi}}(t), \sqrt{\boldsymbol{\gamma}}}\|_2$ that takes advantage of the presence of the vector $\boldsymbol{\gamma}$. It is not hard to see that the L^2 norm of the matrix $(\text{diag}(\hat{\boldsymbol{\pi}}) - \hat{\boldsymbol{\pi}}\hat{\boldsymbol{\pi}}^\top)$ is bounded by $\max_{j \in [q]} \hat{\pi}_j$. The following result can be seen as a generalisation of this fact, which is however significantly more involved to prove. The proof is given in Section 4.

² For a square matrix \mathbf{M} , we use $\|\mathbf{M}\|_2$ to denote its L^2 norm, i.e., $\|\mathbf{M}\|_2 = \max_{\|\mathbf{x}\|_2=1} \|\mathbf{M}\mathbf{x}\|_2$. A fact that will be useful later is that $\|\mathbf{M}\|_2 = \max_{\|\mathbf{x}\|_2=1} \|\mathbf{x}^\top \mathbf{M}\|_2$, even for non-symmetric matrices \mathbf{M} .

³ For a vector \mathbf{x} , $\text{diag}(\mathbf{x})$ denotes the diagonal matrix with the entries of \mathbf{x} on the diagonal.

► **Lemma 11.** *Let q be a positive integer, $\hat{\pi}$ be a q -dimensional probability vector and $\hat{\gamma}$ be a q -dimensional vector with non-negative entries which are all bounded by 1. Then, the L^2 norm of the matrix $\mathbf{M}_{\hat{\pi}, \hat{\gamma}} = (\text{diag}(\hat{\pi}) - \hat{\pi}\hat{\pi}^\top)\text{diag}(\hat{\gamma})$ satisfies*

$$\|\mathbf{M}_{\hat{\pi}, \hat{\gamma}}\|_2 \leq \frac{1}{2} \max_{j \in [q]} \hat{\pi}_j (1 + (\hat{\gamma}_j)^2),$$

where $\{\hat{\pi}_j\}_{j \in [q]}, \{\hat{\gamma}_j\}_{j \in [q]}$ are the entries of $\hat{\pi}, \hat{\gamma}$, respectively.

The final component of our proof is to utilize the bound in Lemma 11 to derive an upper bound on the norm of the matrix $\mathbf{M}_{\hat{\pi}(t), \sqrt{\gamma}}$ appearing in Lemma 10. To prove Theorem 4, we roughly need to show that the norm is bounded by $1/|D|$. We show that this is indeed the case in Section 6.

► **Lemma 12.** *There exist absolute constants $\beta > 0$ and $K' \in (0, 1)$ such that the following holds for all positive integers q, d satisfying $q \geq 1.59d + \beta$.*

Let $\gamma, \hat{\pi}(t)$ be the q -dimensional vectors of Definitions 7 and 8, respectively. Then, for all $t \in [0, 1]$ and all colors $k \in [q]$, it holds that

$$\frac{1}{2} \hat{\pi}_k(t) (1 + \gamma_k) < K' / |D|,$$

where D is the set of non-frozen children of ρ under η and η' .

Assuming Lemmas 10, 11 and 12 for now, we next conclude the proof of Theorem 4.

Proof of Theorem 4. Let $U' := (1 + K')/2$ where $K' \in (0, 1)$ is the constant in Lemma 12. Let $\beta > 0$ be a sufficiently large constant so that, for all $q \geq 1.59d + \beta$, the conclusion of Lemma 12 applies and $\frac{1}{1 - \frac{1}{q-d}} K' < U'$. We will show that

$$\|\pi - \pi'\|_2^2 \leq U \max_{i \in [d]} \|\pi_i - \pi_{i'}\|_2^2, \text{ with } U := (U')^2. \quad (4)$$

Indeed, by Lemmas 10, 11 and 12, we have that

$$\|\pi - \pi'\|_2^2 \leq \frac{U}{|D|} \sum_{i \in [d]} \|\pi_i - \pi_{i'}\|_2^2.$$

Note that an index $i \notin D$ corresponds to a frozen child v_i and therefore $\pi_i = \pi_{i'}$ for all $i \notin D$ and hence

$$\frac{1}{|D|} \sum_{i \in [d]} \|\pi_i - \pi_{i'}\|_2^2 \leq \max_{i \in [d]} \|\pi_i - \pi_{i'}\|_2^2,$$

proving (4). This completes the proof of Theorem 4. ◀

4 Bound on the matrix norm: proof of Lemma 11

In this section, we prove Lemma 11.

Proof of Lemma 11. For this proof, it will be convenient to simplify notation and use π instead of $\hat{\pi}$ and γ instead of $\hat{\gamma}$, so that $\mathbf{M}_{\hat{\pi}, \hat{\gamma}}$ becomes $(\text{diag}(\pi) - \pi\pi^\top)\text{diag}(\gamma)$. Let $C := \frac{1}{2} \max_{j \in [q]} \pi_j (1 + \gamma_j^2)$. We will establish that $\|\mathbf{M}_{\pi, \gamma}\|_2 \leq C$ by showing that for an arbitrary q -dimensional vector \mathbf{x} it holds that

$$\|\mathbf{x}^\top \mathbf{M}_{\pi, \gamma}\|_2^2 \leq C^2 \|\mathbf{x}\|_2^2. \quad (5)$$

48:8 Improved Strong Spatial Mixing for Colorings on Trees

We will focus on proving (5) in the case where the entries of the vector γ are all nonnegative and strictly less than one; the case where some of the entries of γ are equal to 1 follows from the continuity of (5) with respect to γ .

So, assume that $\gamma_j \in [0, 1)$ for all $j \in [q]$. Observe that

$$\|\mathbf{x}^\top \mathbf{M}_{\pi, \gamma}\|_2^2 = \sum_{j \in [q]} \pi_j^2 \gamma_j^2 (x_j - w)^2 \text{ where } w := \sum_{j \in [q]} \pi_j x_j.$$

Let $y_j = x_j - w$ for $j \in [q]$. Since π is a probability vector, we have

$$\sum_{j \in [q]} \pi_j y_j = 0.$$

Moreover, we can rewrite (5) as

$$\sum_{j \in [q]} \frac{\pi_j^2 \gamma_j^2}{C^2} y_j^2 \leq \sum_{j \in [q]} (y_j + w)^2. \quad (6)$$

Note that the function $f(z) = \sum_{j \in [q]} (y_j + z)^2$ achieves its minimum for $z^* = -\frac{1}{q} \sum_{j \in [q]} y_j$ and $f(z^*) = \sum_{j \in [q]} y_j^2 - \frac{1}{q} (\sum_{j \in [q]} y_j)^2$. Hence, to prove (6) (and therefore (5)), it suffices to show that

$$\left(\sum_{j \in [q]} y_j \right)^2 \leq q \sum_{j \in [q]} \frac{y_j^2}{A_j}, \text{ where } A_j := \frac{C^2}{C^2 - \pi_j^2 \gamma_j^2} \quad (7)$$

Note that the A_j 's are well-defined and greater than 1 for all $j \in [q]$ by our assumption that $\gamma_j \in [0, 1)$, cf. the argument below (5). Using that $\sum_{j \in [q]} \pi_j y_j = 0$, we therefore obtain that (7) is equivalent to

$$\left(\sum_{j \in [q]} y_j (1 + t\pi_j) \right)^2 \leq q \sum_{j \in [q]} \frac{y_j^2}{A_j}, \text{ where } A_j := \frac{C^2}{C^2 - \pi_j^2 \gamma_j^2}, \quad (8)$$

for any real number t – we will specify t soon (cf. the upcoming (10)). In particular, by the Cauchy-Schwarz inequality, we have

$$\left(\sum_{j \in [q]} y_j (1 + t\pi_j) \right)^2 \leq \sum_{j \in [q]} \frac{y_j^2}{A_j} \sum_{j \in [q]} A_j (1 + t\pi_j)^2,$$

so (8) and hence (7) will follow if we find t such that

$$\sum_{j \in [q]} A_j (1 + t\pi_j)^2 \leq q. \quad (9)$$

We will choose t to minimise the l.h.s. in (9), i.e., set

$$t := -\frac{\sum_{j \in [q]} A_j \pi_j}{\sum_{j \in [q]} A_j \pi_j^2}, \text{ so that } \sum_{j \in [q]} A_j (1 + t\pi_j)^2 = \sum_{j \in [q]} A_j - \frac{(\sum_{j \in [q]} A_j \pi_j)^2}{\sum_{j \in [q]} A_j \pi_j^2}. \quad (10)$$

Therefore, for this choice of t , (9) becomes

$$\sum_{j \in [q]} (A_j - 1) \sum_{j \in [q]} A_j \pi_j^2 \leq \left(\sum_{j \in [q]} A_j \pi_j \right)^2. \quad (11)$$

Using that $A_j = \frac{C^2}{C^2 - \pi_j^2 \gamma_j^2}$, (11) is equivalent to (note the division by C^2 of both sides)

$$\sum_{j \in [q]} \frac{\pi_j^2 \gamma_j^2}{C^2 - \pi_j^2 \gamma_j^2} \sum_{j \in [q]} \frac{\pi_j^2}{C^2 - \pi_j^2 \gamma_j^2} \leq \left(\sum_{j \in [q]} \frac{C \pi_j}{C^2 - \pi_j^2 \gamma_j^2} \right)^2. \quad (12)$$

We next establish (12). We can upper bound the l.h.s. of (12) using the inequality $ab \leq \left(\frac{a+b}{2}\right)^2$, which gives that

$$\sum_{j \in [q]} \frac{\pi_j^2 \gamma_j^2}{C^2 - \pi_j^2 \gamma_j^2} \sum_{j \in [q]} \frac{\pi_j^2}{C^2 - \pi_j^2 \gamma_j^2} \leq \left(\sum_{j \in [q]} \frac{\pi_j^2 (1 + \gamma_j^2)}{2(C^2 - \pi_j^2 \gamma_j^2)} \right)^2.$$

So, to prove (12), it suffices to show that for each $i \in [q]$, it holds that

$$\frac{\pi_j^2 (1 + \gamma_j^2)}{2(C^2 - \pi_j^2 \gamma_j^2)} \leq \frac{C \pi_j}{C^2 - \pi_j^2 \gamma_j^2}$$

which is indeed true, since $C \geq \frac{1}{2} \pi_j (1 + \gamma_j^2)$ for all $i \in [q]$ by the definition of C .

This proves (12), which in turn establishes (8) for the choice of t in (10). This yields (7) and hence (5) as well, finishing the proof of Lemma 11. \blacktriangleleft

5 Gradient analysis with blocked colors: proof of Lemma 10

In this section, we prove Lemma 10.

Proof of Lemma 10. For $i \in [d]$ and $j \in [q]$, let $F_{c,j}^{(i)}(\mathbf{x})$ be the partial derivative $\frac{\partial f_c}{\partial x_{i,j}}$ viewed as a function of the “concatenated” vector $\mathbf{x} = (\mathbf{x}_1, \dots, \mathbf{x}_d)$. Note that, whenever $x_{i,j} \neq 1$, we have that

$$\begin{aligned} F_{c,j}^{(i)}(\mathbf{x}) &= -\frac{f_c(\mathbf{x}_1, \dots, \mathbf{x}_d) - (f_c(\mathbf{x}_1, \dots, \mathbf{x}_d))^2}{1 - x_{i,j}} \text{ if } j = c, \\ F_{c,j}^{(i)}(\mathbf{x}) &= \frac{f_c(\mathbf{x}_1, \dots, \mathbf{x}_d) f_j(\mathbf{x}_1, \dots, \mathbf{x}_d)}{1 - x_{i,j}} \text{ if } j \neq c. \end{aligned} \quad (13)$$

As mentioned earlier, we will interpolate between $\boldsymbol{\pi}$ and $\boldsymbol{\pi}'$ by interpolating along the straight-line segment connecting $(\boldsymbol{\pi}_1, \dots, \boldsymbol{\pi}_d)$ and $(\boldsymbol{\pi}'_1, \dots, \boldsymbol{\pi}'_d)$. In particular, for $t \in [0, 1]$, let $\hat{\pi}_c(t)$ denote the c -th entry of the vector $\hat{\boldsymbol{\pi}}(t)$ defined in the statement of the lemma. Then, we have that

$$\hat{\pi}_c(t) = f_c(\mathbf{z}(t)), \text{ where } \mathbf{z}(t) \text{ is the vector } (t\boldsymbol{\pi}_1 + (1-t)\boldsymbol{\pi}'_1, \dots, t\boldsymbol{\pi}_d + (1-t)\boldsymbol{\pi}'_d). \quad (14)$$

We will use $z_{i,j}(t)$ to denote the j -th entry of the i -th vector in $\mathbf{z}(t)$, i.e., $z_{i,j}(t) = t\pi_{i,j} + (1-t)\pi'_{i,j}$.

Let D be the set of indices i such that v_i is not frozen under η and η' (cf. Observation 6). Observe that, for all $i \notin D$ and $c, j \in [q]$, we have that $z_{i,j}(t) = \pi_{i,j} = \pi'_{i,j}$ for $t \in [0, 1]$. Moreover, for $i \in D$ and $j \in [q]$ we have that $\pi_{i,j}, \pi'_{i,j} \leq 1/(q-d)$ (since the child v_i has at least $q-d$ available colors in the subtree T_i) and hence

$$0 \leq z_{i,j}(t) \leq 1/(q-d). \quad (15)$$

Since $z_{i,j}(t) \neq 1$ for $i \in D$ and $j \in [q]$, it follows that

$$\frac{d\hat{\pi}_c}{dt} = \sum_{i \in D} \sum_{j=1}^q F_{c,j}^{(i)}(\mathbf{z}(t)) (\pi_{i,j} - \pi'_{i,j}).$$

48:10 Improved Strong Spatial Mixing for Colorings on Trees

Using (3), we therefore have that

$$\begin{aligned} (\pi_c - \pi'_c)^2 &= (\hat{\pi}_c(1) - \hat{\pi}_c(0))^2 = \left(\int_0^1 \frac{d\hat{\pi}_c}{dt} dt \right)^2 = \left(\int_0^1 \sum_{i \in D} \sum_{j=1}^q F_{c,j}^{(i)}(\mathbf{z}(t)) (\pi_{i,j} - \pi'_{i,j}) dt \right)^2 \\ &\leq \int_0^1 \left(\sum_{i \in D} \sum_{j=1}^q F_{c,j}^{(i)}(\mathbf{z}(t)) (\pi_{i,j} - \pi'_{i,j}) \right)^2 dt, \end{aligned}$$

where the last inequality follows by applying the Cauchy-Schwarz inequality for integrals. By summing over all colors $c \in [q]$, we obtain

$$\|\boldsymbol{\pi} - \boldsymbol{\pi}'\|_2^2 \leq \int_0^1 \sum_{c=1}^q \left(\sum_{i \in D} \sum_{j=1}^q F_{c,j}^{(i)}(\mathbf{z}(t)) (\pi_{i,j} - \pi'_{i,j}) \right)^2 dt. \quad (16)$$

To simplify the r.h.s. of (16), we first note that, by (13) and (14), we have

$$F_{c,j}^{(i)}(\mathbf{z}(t)) = \frac{A_{c,j}(t)}{1 - z_{i,j}(t)} \text{ where } A_{c,j} := \begin{cases} (\hat{\pi}_c(t))^2 - \hat{\pi}_c(t), & \text{if } j = c, \\ \hat{\pi}_c(t)\hat{\pi}_j(t), & \text{if } j \neq c \end{cases} \quad (17)$$

Moreover, for $j \in [q]$, set

$$u_j(t) = \frac{1}{|D|\gamma_j} \sum_{i \in D} \frac{\pi_{i,j} - \pi'_{i,j}}{1 - z_{i,j}(t)} \text{ if } \gamma_j > 0, \text{ else set } u_j(t) = 0. \quad (18)$$

Note that if color j is blocked for the child v_i we have that $\pi_{i,j} - \pi'_{i,j} = 0$, so using the power mean inequality we have that

$$\gamma_j (u_j(t))^2 \leq \frac{1}{|D|} \sum_{i \in D} \left(\frac{\pi_{i,j} - \pi'_{i,j}}{1 - z_{i,j}(t)} \right)^2. \quad (19)$$

Then, for $c \in [q]$, we have that

$$\sum_{i \in D} \sum_{j=1}^q F_{c,j}^{(i)}(\mathbf{z}(t)) (\pi_{i,j} - \pi'_{i,j}) = \sum_{j=1}^q A_{c,j}(t) \sum_{i \in D} \frac{\pi_{i,j} - \pi'_{i,j}}{1 - z_{i,j}(t)} = |D| \sum_{j=1}^q A_{c,j}(t) \gamma_j u_j(t), \quad (20)$$

where the last equality follows from (18) and observing that if $\gamma_j = 0$ then $\pi_{i,j} - \pi'_{i,j} = 0$ for all $i \in D$. Note that the (c, q) -entry of $\mathbf{M}_{\hat{\boldsymbol{\pi}}(t), \sqrt{\boldsymbol{\gamma}}}$ is exactly $-A_{c,j}(t)\sqrt{\gamma_j}$ (cf. (17) and Definition 9) and hence, using (20), we can write the integrand in the r.h.s. of (16) as

$$\sum_{c=1}^q \left(\sum_{i \in D} \sum_{j=1}^q F_{c,j}^{(i)}(\mathbf{z}(t)) (\pi_{i,j} - \pi'_{i,j}) \right)^2 = |D|^2 \|\mathbf{M}_{\hat{\boldsymbol{\pi}}(t), \sqrt{\boldsymbol{\gamma}}} \mathbf{u}(t)\|_2^2, \quad (21)$$

where, for $t \in [0, 1]$, $\mathbf{u}(t)$ is the q -dimensional vector with entries $\{\sqrt{\gamma_j} u_j(t)\}_{j \in [q]}$. Let

$$W := \max_{t \in [0,1]} \|\mathbf{M}_{\hat{\boldsymbol{\pi}}(t), \sqrt{\boldsymbol{\gamma}}}\|_2, \text{ so that } K = \frac{W}{1 - \frac{1}{q-d}}.$$

Then, for $t \in [0, 1]$, we have that

$$\begin{aligned} \|\mathbf{M}_{\hat{\boldsymbol{\pi}}(t), \sqrt{\boldsymbol{\gamma}}} \mathbf{u}(t)\|_2^2 &\leq W^2 \|\mathbf{u}(t)\|_2^2 = W^2 \sum_{j \in [q]} \gamma_j (u_j(t))^2 \leq \frac{W^2}{|D|} \sum_{j \in [q]} \sum_{i \in D} \left\| \frac{\pi_{i,j} - \pi'_{i,j}}{1 - z_{i,j}(t)} \right\|_2^2 \\ &\leq \frac{K^2}{|D|} \sum_{j \in [q]} \sum_{i \in D} \|\pi_{i,j} - \pi'_{i,j}\|_2^2 = \frac{K^2}{|D|} \sum_{i \in [d]} \|\boldsymbol{\pi}_i - \boldsymbol{\pi}'_i\|_2^2, \end{aligned} \quad (22)$$

where the first inequality is by definition of the norm, the second inequality follows from (19), the third inequality follows from $0 \leq z_{i,j}(t) \leq 1/(q-d)$, and the last equality follows from the fact that for $i \notin D$ we have that $\pi_i = \pi'_i$. Combining (16), (21) and (22), we obtain that

$$\|\pi - \pi'\|_2^2 \leq |D|K^2 \sum_{i \in [d]} \|\pi_i - \pi'_i\|_2^2.$$

This finishes the proof of Lemma 10. \blacktriangleleft

6 Bounds on the marginals: proof of Lemma 12

In this section, we prove Lemma 12. We begin with the following lemma.

► **Lemma 13.** *Let q, d, h be positive integers so that $q \geq d+1$ and $h \geq 1$. Let $T = \mathbb{T}_{d,h,\rho}$ be the d -ary tree with height h rooted at ρ , Λ be a subset of the vertices of T such that $\rho \notin \Lambda$, and $\eta : \Lambda \rightarrow [q]$ be an extendible assignment of T . Then, for all colors $k \in [q]$ that are available for ρ under η , it holds that*

$$\mu_T(\sigma_\rho = k \mid \sigma_\Lambda = \eta) \geq \frac{(1 - \frac{1}{q-d})^d}{d + (q-d)(1 - \frac{1}{q-d})^d}.$$

Proof. Let $Q \subseteq [q]$ be the set of all colors that are available for ρ under η and let $k \in Q$. Let v_1, \dots, v_d be the children of ρ in T and let $D = \{i \in [d] \mid v_i \notin \Lambda\}$ be the indices of the children of ρ that do not belong to Λ .

For $i \in [d]$, let $T_i = (V_i, E_i)$ be the subtree of T rooted at v_i which consists of all descendants of v_i in T (together with v_i itself). Further, for a color $j \in [q]$, let

$$x_{i,j} = \mu_{T_i}(\sigma_{v_i} = j \mid \sigma_{\Lambda \cap V_i} = \eta_{\Lambda \cap V_i}),$$

i.e., $x_{i,j}$ is the marginal probability that v_i takes the color j at v_i in μ_{T_i} with boundary condition $\eta_{\Lambda \cap V_i}$. Note that

$$0 \leq x_{i,j} \leq \frac{1}{q-d} \text{ for all } i \in D \text{ and } j \in [q], \quad \sum_{j \in [q]} x_{i,j} = 1 \text{ for all } i \in [q]. \quad (23)$$

Using the tree recursion (2) and ignoring summands that are 0 or factors that are equal to 1, the marginal $\mu_T(\sigma_\rho = k \mid \sigma_\Lambda = \eta)$ is expressed in terms of $x_{i,j}$ as follows:

$$\mu_T(\sigma_\rho = k \mid \sigma_\Lambda = \eta) = \frac{\prod_{i \in D} (1 - x_{i,k})}{\sum_{j \in Q} \prod_{i \in D} (1 - x_{i,j})}. \quad (24)$$

We prove the lemma by deriving an appropriate lower bound on the quantity at the r.h.s. of (24) subject to the constraint in (23). For the numerator in (24), we have that

$$\prod_{i \in D} (1 - x_{i,k}) \geq \left(1 - \frac{1}{q-d}\right)^{|D|}. \quad (25)$$

For the denominator we are going to show the following:

$$\sum_{j \in Q} \prod_{i \in D} (1 - x_{i,j}) \leq d + (q-d) \left(1 - \frac{1}{q-d}\right)^{|D|}. \quad (26)$$

48:12 Improved Strong Spatial Mixing for Colorings on Trees

Before showing that (26) is indeed true, note that the lemma follows by plugging (25), (26) into (24), yielding

$$\mu_T(\sigma_\rho = k \mid \sigma_\Lambda = \eta) \geq \frac{\left(1 - \frac{1}{q-d}\right)^{|D|}}{d + (q-d) \left(1 - \frac{1}{q-d}\right)^{|D|}} \geq \frac{\left(1 - \frac{1}{q-d}\right)^d}{d + (q-d) \left(1 - \frac{1}{q-d}\right)^d},$$

where the last inequality follows by noting that the ratio in the middle is decreasing in $|D|$ and $|D| \leq d$.

We now proceed with the proof of (26). First, we have the simple bound

$$\sum_{j \in Q} \prod_{i \in D} (1 - x_{i,j}) \leq \sum_{j \in [q]} \prod_{i \in D} (1 - x_{i,j}). \quad (27)$$

For $j \in [q]$, let $x_j = \frac{1}{|D|} \sum_{i \in D} x_{i,j}$ and note that (x_1, \dots, x_q) is a probability vector whose entries are in $[0, 1/(q-d)]$. By the AM-GM inequality, we can bound the r.h.s. of (27) by

$$\sum_{j \in [q]} \prod_{i \in D} (1 - x_{i,j}) \leq \sum_{j \in [q]} (1 - x_j)^{|D|}. \quad (28)$$

It remains to observe that the function $f(\mathbf{z}) = \sum_{j \in [q]} (1 - z_j)^{|D|}$ is convex over the space of probability vectors $\mathbf{z} = (z_1, \dots, z_q)$ whose entries are in $[0, 1/(q-d)]$, and hence f attains its maximum at the extreme points of the space, which are given by (the permutations of) the probability vector whose first d entries are equal to zero and the rest are equal to $1/(q-d)$. It follows that

$$\sum_{j \in [q]} (1 - x_j)^{|D|} \leq d + (q-d) \left(1 - \frac{1}{q-d}\right)^{|D|}. \quad (29)$$

Combining (27), (28) and (29) yields (26), thus concluding the proof of Lemma 13. \blacktriangleleft

We are now ready to prove Lemma 12.

Proof of Lemma 12. For convenience, let $r = 1.59$, so that $q/d \geq r$. We will use that r satisfies

$$C := \frac{1}{r} \exp\left(\frac{1}{r}\right) \exp\left(-\frac{1}{r-1 + \exp\left(\frac{1}{r-1}\right)}\right) < 1. \quad (30)$$

We will show the result with the constant $K' = (1 + C)/2$. For the rest of this proof, we will focus on the case $q \in [1.59d + \beta, 2.01d]$, for some large constant $\beta > 0$ (when $q > 2.01d$ the desired bound follows rather crudely, see Footnote 4 below for details).

Recall that v_1, \dots, v_d are the children of ρ in T and D is the set of (indices of the) non-frozen children of the root ρ . Let $Q \subseteq [q]$ be the set of all colors that are available for ρ under η ; since at most $d - |D|$ colors can be blocked for ρ , we have that

$$|Q| \geq q - (d - |D|). \quad (31)$$

For $i \in [d]$, let $T_i = (V_i, E_i)$ be the subtree of T rooted at v_i which consists of all descendants of v_i in T (together with v_i itself). Further, for a color $j \in [q]$, recall that

$$\begin{aligned} \pi_{i,j} &= \mu_{T_i}(\sigma_{v_i} = j \mid \sigma_{\Lambda \cap V_i} = \eta_{\Lambda \cap V_i}), \\ \pi'_{i,j} &= \mu_{T_i}(\sigma_{v_i} = j \mid \sigma_{\Lambda \cap V_i} = \eta'_{\Lambda \cap V_i}), \end{aligned} \quad (32)$$

i.e., $\pi_{i,j}$ is the marginal probability that v_i takes the color j at v_i in μ_{T_i} with boundary condition $\eta_{\Lambda \cap V_i}$. For a non-frozen child v_i (i.e., $i \in D$), note that, if color j is available for v_i (in T_i), then we have from Lemma 13 the bounds

$$L \leq \pi_{i,j}, \pi_{i,j}, \text{ where } L = \frac{\left(1 - \frac{1}{q-d}\right)^d}{d + (q-d)\left(1 - \frac{1}{q-d}\right)^d}. \quad (33)$$

Another useful bound to observe for later is that

$$dL < 1/3 \text{ for all } d \geq 2.$$

Consider arbitrary $k \in Q$. For $t \in [0, 1]$, let $\mathbf{z}(t)$ be the vector $(t\pi_1 + (1-t)\pi'_1, \dots, t\pi_d + (1-t)\pi'_d)$. Using the tree recursion (2) and ignoring summands that are 0 or factors that are equal to 1, we obtain

$$\hat{\pi}_k(t) = \frac{\prod_{i \in D} (1 - z_{i,k}(t))}{\sum_{j \in Q} \prod_{i \in D} (1 - z_{i,j}(t))}. \quad (34)$$

Recall, our goal is to show that $\frac{1}{2}\hat{\pi}_k(t)(1 + \gamma_k) < K'/|D|$ for all $t \in [0, 1]$, where $\gamma_k \in [0, 1]$ is the fraction of non-frozen children that have color k available. ⁴Note that, if color j is available for the child v_i , (33) gives that

$$L \leq z_{i,j}(t) \text{ for } t \in [0, 1],$$

so, using the fact that the color k is available for $|D|\gamma_k$ non-frozen children, we obtain that the numerator of (34) is bounded by

$$\prod_{i \in D} (1 - z_{i,k}(t)) \leq (1 - L)^{|D|\gamma_k} \leq \exp(-L|D|\gamma_k), \quad (35)$$

whereas the denominator, using the AM-GM inequality analogously to [7, Lemma 2.1], by

$$\sum_{j \in Q} \prod_{i \in D} (1 - z_{i,j}(t)) \geq q \exp(-|D|/q) - (d - |D|). \quad (36)$$

From (34), (35), and (36), it follows that $\hat{\pi}_k(t) \leq \frac{\exp(-L|D|\gamma_k)}{q \exp(-|D|/q) - (d - |D|)}$. Therefore, the lemma will follow by showing that

$$\frac{|D| \exp(-|D|L\gamma_k)}{q \exp(-|D|/q) - (d - |D|)} (1 + \gamma_k) < 2K'. \quad (37)$$

Note that the function $h(x) = (1 + x) \exp(-dLx)$ is increasing when $x \in [0, 1]$, since

$$h'(x) = \exp(-dLx)(1 - dL(1 + x)) \geq \exp(-dLx)(1 - 2dL) > 0.$$

Therefore, to prove (37), it suffices to show that

$$\frac{|D| \exp(-|D|L)}{q \exp(-|D|/q) - (d - |D|)} < K', \text{ or equivalently that } f(|D|) > 0 \quad (38)$$

⁴ For $q > 2.01d$, we have from (34) and (31) that $\hat{\pi}_k(t) \leq \frac{1}{|Q|-|D|} \leq \frac{1}{q-d} < \frac{1}{1.01d} \leq K'/|D|$, yielding the desired inequality.

48:14 Improved Strong Spatial Mixing for Colorings on Trees

where $f(x) := K'(q \exp(-x/q) - d + x) - x \exp(-Lx)$ for $x \in [0, d]$. We claim that $f(x)$ is decreasing in x . We have

$$f'(x) = K' - K' \exp(-x/q) - \exp(-Lx)(1 - Lx)$$

which is maximised for $x = d$. In particular,

$$\begin{aligned} f'(x) &\leq f'(d) = K' - K' \exp(-d/q) - \exp(-dL)(1 - dL) \\ &\leq K' - K' \exp(-1/r) - \exp(-1/3)(1 - 1/3) \leq 0, \end{aligned}$$

where the second to last inequality follows from the fact that $dL < 1/3$ and the last inequality using that $K' < 1$. For $|D| = d$, (38) becomes

$$\frac{d \exp(-dL)}{q \exp(-d/q)} < K'. \quad (39)$$

Now, we have that

$$dL \geq \frac{1}{r - 1 + \exp\left(\frac{d}{(r-1)d-1}\right)}.$$

Therefore, by choosing β large enough, we can ensure that

$$\frac{d \exp(-dL)}{q \exp(-d/q)} < \frac{1 + C}{2} = K',$$

where C is the constant in (30). This proves (39) and therefore concludes the proof of Lemma 12. \blacktriangleleft

7 Proof of Theorem 3

Finally, utilizing Theorem 4, we give the proof of Theorem 3.

Proof of Theorem 3. From Theorem 4, we know that there exist constants $\beta > 0$ and $U \in (0, 1)$ such that for all $q \geq 1.59d + \beta$ the conclusion of Theorem 4 applies. Note that Theorem 4 applies to the d -ary tree rather than the $(d+1)$ -regular tree but these trees differ only at the degree of the root. To account for it, we will assume that $q \geq 1.59(d+1) + \beta$, i.e., prove Theorem 3 with constant $\beta' = \beta + 1.59$. Consider the function ζ given by $\zeta(\ell) = 2U^{\ell-2}$ for $\ell \geq 0$ and note that ζ is exponentially decaying. We will show that the q -coloring model has strong spatial mixing on the $(d+1)$ -regular tree with decay rate ζ .

We first show by induction on h that, for the tree $T = \hat{\mathbb{T}}_{d+1, h, \rho}$ (that is, the $(d+1)$ -ary tree with height h rooted at ρ), for any subset Λ of vertices of T and arbitrary extendible assignments $\eta, \eta' : \Lambda \rightarrow [q]$ of T , it holds that

$$\|\pi_{T, \rho, \eta} - \pi_{T, \rho, \eta'}\|_2^2 \leq \zeta(\text{dist}(\rho, \Delta)), \quad (40)$$

where $\Delta \subseteq \Lambda$ is the set of vertices where η and η' disagree. The base cases $h = 0, 1, 2$ are trivial so assume $h \geq 3$ in what follows. Let $\ell = \text{dist}(\rho, \Delta)$. Once again, (40) is trivial when $\ell \leq 2$, so assume $\ell \geq 3$ in what follows. Let v_1, \dots, v_{d+1} be the children of ρ and, for $i \in [d+1]$, let $T_i = (V_i, E_i)$ be the subtree of T rooted at v_i which consists of all descendants of v_i in T . Further, let $\pi_i = \pi_{T_i, v_i, \eta(\Lambda \cap V_i)}$, $\pi'_i = \pi_{T_i, v_i, \eta'(\Lambda \cap V_i)}$. Then, by Theorem 4 and since $q \geq 1.59(d+1) + \beta$, we have that

$$\|\pi_{T, \rho, \eta} - \pi_{T, \rho, \eta'}\|_2^2 \leq U \max_{i \in [d+1]} \|\pi_i - \pi'_i\|_2^2. \quad (41)$$

For $i \in [d+1]$, since T_i is isomorphic to $\hat{\mathbb{T}}_{d+1, h-1, \rho}$ we have by the induction hypothesis that

$$\|\pi_i - \pi_{i'}\|_2^2 \leq \zeta(\ell - 1).$$

Combining this with (41) and the fact that $\zeta(\ell) = U\zeta(\ell - 1)$ yields (40), completing the induction and therefore that strong spatial mixing holds on T with decay rate ζ .

Now, let $T = (V, E)$ be a finite subtree of the $(d+1)$ -regular tree, v be an arbitrary vertex of T , Λ be a subset of vertices of T and $\eta, \eta' : \Lambda \rightarrow [q]$ be arbitrary extendible assignments of T . Then, we can view T as a subgraph of $T_v = \hat{\mathbb{T}}_{d+1, h, v}$ for some appropriate height h . It also holds that (see, for example, [10, Lemma 25])

$$\|\pi_{T, v, \eta} - \pi_{T, v, \eta'}\|_2 = \|\pi_{T_v, v, \eta} - \pi_{T_v, v, \eta'}\|_2.$$

Therefore, from (40) (applied to the tree T_v) we obtain that

$$\|\pi_{T, v, \eta} - \pi_{T, v, \eta'}\|_2^2 \leq \zeta(\text{dist}(v, \Delta)),$$

where $\Delta \subseteq \Lambda$ is the set of vertices where η and η' disagree.

This completes the proof of Theorem 3. ◀

References

- 1 A. Barvinok. *Combinatorics and Complexity of Partition Functions*. Algorithms and Combinatorics. Springer International Publishing, 2017.
- 2 V. Beffara and H. Duminil-Copin. The self-dual point of the two-dimensional random-cluster model is critical for $q \geq 1$. *Probability Theory and Related Fields*, 153(3):511–542, 2012.
- 3 A. Blanca, P. Caputo, A. Sinclair, and E. Vigoda. Spatial Mixing and Non-local Markov Chains. In *Proceedings of the Twenty-Ninth Annual ACM-SIAM Symposium on Discrete Algorithms*, SODA '18, pages 1965–1980, 2018.
- 4 A. Blanca and A. Sinclair. Random-cluster dynamics in \mathbb{Z}^2 . *Probability Theory and Related Fields*, 168(3-4):821–847, 2017.
- 5 G. R. Brightwell and P. Winkler. Random Colorings of a Cayley Tree. In *Contemporary Combinatorics*, pages 247–276, 2002.
- 6 F. Cesi. Quasi-factorization of the entropy and logarithmic Sobolev inequalities for Gibbs random fields. *Probability Theory and Related Fields*, 120(4):569–584, 2001.
- 7 M. Dyer and A. Frieze. Randomly coloring graphs with lower bounds on girth and maximum degree. *Random Structures & Algorithms*, 23(2):167–179, 2003.
- 8 M. Dyer, A. Frieze, T. P. Hayes, and E. Vigoda. Randomly coloring constant degree graphs. *Random Structures & Algorithms*, 43(2):181–200, 2013.
- 9 M. Dyer, A. Sinclair, E. Vigoda, and D. Weitz. Mixing in time and space for lattice spin systems: A combinatorial view. *Random Structures & Algorithms*, 24(4):461–479, 2004.
- 10 C. Efthymiou. A Simple Algorithm for Sampling Colorings of $G(n, d/n)$ Up to The Gibbs Uniqueness Threshold. *SIAM Journal on Computing*, 45(6):2087–2116, 2016.
- 11 A. Galanis, L. A. Goldberg, and K. Yang. Uniqueness for the 3-state antiferromagnetic Potts model on the tree. *Electron. J. Probab.*, 23, 2018.
- 12 D. Gamarnik, D. Katz, and S. Misra. Strong spatial mixing of list coloring of graphs. *Random Structures & Algorithms*, 46(4):599–613, 2015.
- 13 Q. Ge and D. Štefankovič. Strong spatial mixing of q -colorings on Bethe lattices. *CoRR*, abs/1102.2886, 2011. [arXiv:1102.2886](https://arxiv.org/abs/1102.2886).
- 14 L. A. Goldberg, R. Martin, and M. Paterson. Strong Spatial Mixing with Fewer Colors for Lattice Graphs. *SIAM Journal on Computing*, 35(2):486–517, 2005.
- 15 T. P. Hayes. Local uniformity properties for Glauber dynamics on graph colorings. *Random Structures & Algorithms*, 43(2):139–180, 2013.

- 16 J. Jonasson. Uniqueness of uniform random colorings of regular trees. *Statistics & Probability Letters*, 57(3):243–248, 2002.
- 17 F. P. Kelly. Stochastic Models of Computer Communication Systems. *Journal of the Royal Statistical Society. Series B (Methodological)*, 47(3):379–395, 1985.
- 18 L. Li, P. Lu, and Y. Yin. Correlation Decay Up to Uniqueness in Spin Systems. In *Proceedings of the Twenty-fourth Annual ACM-SIAM Symposium on Discrete Algorithms*, SODA '13, pages 67–84, 2013.
- 19 J. Liu and P. Lu. FPTAS for #BIS with degree bounds on one side. In *Proceedings of the Forty-seventh Annual ACM Symposium on Theory of Computing*, STOC '15, pages 549–556, 2015.
- 20 J. Liu, A. Sinclair, and P. Srivastava. A deterministic algorithm for counting colorings with 2Δ colors. *CoRR*, abs/1906.01228, 2019. [arXiv:1906.01228](https://arxiv.org/abs/1906.01228).
- 21 F. Martinelli and E. Olivieri. Approach to equilibrium of Glauber dynamics in the one phase region. I. The attractive case. *Communications in Mathematical Physics*, 161(3):447–486, 1994.
- 22 F. Martinelli and E. Olivieri. Approach to equilibrium of Glauber dynamics in the one phase region. II. The General Case. *Communications in Mathematical Physics*, 161(3):487–514, 1994.
- 23 V. Patel and G. Regts. Deterministic Polynomial-Time Approximation Algorithms for Partition Functions and Graph Polynomials. *SIAM Journal on Computing*, 46(6):1893–1919, 2017.
- 24 J. Pearl. Reverend Bayes on Inference Engines: A Distributed Hierarchical Approach. In *Proceedings of the Second AAAI Conference on Artificial Intelligence*, AAAI'82, pages 133–136, 1982.
- 25 H. Peters and G. Regts. On a Conjecture of Sokal Concerning Roots of the Independence Polynomial. *The Michigan Mathematical Journal*, pages 33–55, 2019.
- 26 D. Weitz. Counting Independent Sets Up to the Tree Threshold. In *Proceedings of the Thirty-eighth Annual ACM Symposium on Theory of Computing*, STOC '06, pages 140–149, 2006.