

Bounded VC-Dimension Implies the Schur-Erdős Conjecture

Jacob Fox

Department of Mathematics, Stanford University, Stanford, CA, USA
jacobfox@stanford.edu

János Pach

Alfréd Rényi Institute of Mathematics, Budapest, Hungary
IST, Vienna, Austria
MIPT, Moscow, Russia
pach@cims.nyu.edu

Andrew Suk

Department of Mathematics, University of California San Diego, La Jolla, CA, USA
asuk@ucsd.edu

Abstract

In 1916, Schur introduced the Ramsey number $r(3; m)$, which is the minimum integer $n > 1$ such that for any m -coloring of the edges of the complete graph K_n , there is a monochromatic copy of K_3 . He showed that $r(3; m) \leq O(m!)$, and a simple construction demonstrates that $r(3; m) \geq 2^{\Omega(m)}$. An old conjecture of Erdős states that $r(3; m) = 2^{\Theta(m)}$. In this note, we prove the conjecture for m -colorings with bounded VC-dimension, that is, for m -colorings with the property that the set system induced by the neighborhoods of the vertices with respect to each color class has bounded VC-dimension.

2012 ACM Subject Classification Mathematics of computing → Combinatoric problems

Keywords and phrases Ramsey theory, VC-dimension, Multicolor Ramsey numbers

Digital Object Identifier 10.4230/LIPIcs.SoCG.2020.46

Funding *Jacob Fox*: Supported by a Packard Fellowship and by NSF award DMS-1855635.

János Pach: Supported by Austrian Science Fund (FWF) grant Z 342-N31 and by the Ministry of Education and Science of the Russian Federation in the framework of MegaGrant no 075-15-2019-1926.

Andrew Suk: Supported by NSF CAREER award DMS-1800746 and an Alfred Sloan Fellowship.

1 Introduction

Given n points and n lines in the plane, their incidence graph is a bipartite graph G that contains no $K_{2,2}$ as a subgraph. By a theorem of Erdős [5] and Kővári-Sós-Turán [15], this implies that the number of incidences between the points and the lines is $O(n^{3/2})$. However, a celebrated theorem of Szemerédi and Trotter [20] states that the actual number of incidences is much smaller, only $O(n^{4/3})$, and this bound is tight. There are many similar examples, where extremal graph theory is applicable, but does not yield optimal results. What is behind this curious phenomenon? In the above and in many other examples, the vertices of G are, or can be associated with, points in a Euclidean space, and the fact whether two vertices are connected by an edge can be determined by evaluating a bounded number of polynomials in the coordinates of the corresponding points. In other words, G is a *semi-algebraic graph* of bounded complexity. As was proved by the authors, in collaboration with Sheffer and Zahl [8], for semi-algebraic graphs, one can explore the *geometric* properties of the polynomial surfaces, including separator theorems and the so-called polynomial method [12], to obtain




© Jacob Fox, János Pach, and Andrew Suk;
licensed under Creative Commons License CC-BY
36th International Symposium on Computational Geometry (SoCG 2020).

Editors: Sergio Cabello and Danny Z. Chen; Article No. 46; pp. 46:1–46:8
Leibniz International Proceedings in Informatics



LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany



much stronger results for the extremal graph-theoretic problems in question. In particular, every $K_{2,2}$ -free semi-algebraic graph of n vertices with the complexity parameters associated with the point-line incidence problem has $O(n^{4/3})$ edges. This implies the Szemerédi-Trotter theorem.

There is a fast growing body of literature demonstrating that many important results in extremal combinatorics can be substantially improved, and several interesting conjectures proved, if we restrict our attention to semi-algebraic graphs and hypergraphs; see, e.g., [1, 7, 9]. It is a major unsolved problem to decide whether this partly algebraic and partly geometric assumption can be relaxed and replaced by a purely combinatorial condition. A natural candidate is that the graph has *bounded Vapnik-Chervonenkis dimension* (in short, VC-dimension). The *VC-dimension of a set system* (hypergraph) \mathcal{F} on the ground set V is the *largest* integer d for which there exists a d -element set $S \subset V$ such that for every subset $B \subset S$, one can find a member $A \in \mathcal{F}$ with $A \cap S = B$. The *VC-dimension of a graph* $G = (V, E)$ is the VC-dimension of the set system formed by the neighborhoods of the vertices, where the neighborhood of $v \in V$ is $N(v) = \{u \in V : uv \in E\}$. The VC-dimension, introduced by Vapnik and Chervonenkis [21], is one of the most useful combinatorial parameters that measures the complexity of graphs and hypergraphs. It proved to be relevant in many branches of pure and applied mathematics, including statistics, logic, learning theory, and real algebraic geometry. It has completely transformed combinatorial and computational geometry after its introduction to the subject by Haussler and Welzl [14] in 1987. But can it be applied to extremal graph theory problems?

At first glance, this looks rather unlikely. Returning to our initial example, it is easy to verify that the Vapnik-Chervonenkis dimension of every $K_{2,2}$ -free graph is at most 2. Therefore, we cannot possibly improve the $O(n^{3/2})$ upper bound on the number of edges of $K_{2,2}$ -free graphs by restricting our attention to graphs of bounded VC-dimension. Yet the goal of the present note is to solve the *Schur-Erdős problem*, one of the oldest open questions in Ramsey theory, by placing this restriction.

To describe the problem, we need some notation. For integers $k \geq 3$ and $m \geq 2$, the *Ramsey number* $r(k; m)$ is the smallest integer n such that any m -coloring of the edges of the complete n -vertex graph contains a monochromatic copy of K_k . For the special case when $k = 3$, Issai Schur [19] showed that

$$\Omega(2^m) \leq r(3; m) \leq O(m!).$$

While the form of the upper bound has remained unchanged over the last century, the lower bound was successively improved. The current record is due to Xiaodong *et al.* [22] who showed that $r(3; m) \geq \Omega(3.199^m)$. It is a major open problem in Ramsey theory to close the gap between the lower and upper bounds for $r(3; m)$. Erdős [3] offered cash prizes for solutions to the following problems.

► **Conjecture 1** (\$100). We have $\lim_{m \rightarrow \infty} (r(3; m))^{1/m} < \infty$.

It was shown by Chung [4] that $r(3; m)$ is supermultiplicative, so that the above limit exists.

► **Problem 2** (\$250). Determine $\lim_{m \rightarrow \infty} (r(3; m))^{1/m}$.

It will be more convenient to work with the *dual VC-dimension*. The *dual* of a set system \mathcal{F} is the set system \mathcal{F}^* obtained by interchanging the roles of V and \mathcal{F} . That is, the ground set of \mathcal{F}^* is \mathcal{F} , and

$$\mathcal{F}^* = \{\{A \in \mathcal{F} : v \in A\} : v \in V\}.$$

We say that \mathcal{F} has *dual VC-dimension* d if \mathcal{F}^* has VC-dimension d . Notice that $(\mathcal{F}^*)^* = \mathcal{F}$, and it is known that if \mathcal{F} has VC-dimension d , then \mathcal{F}^* has VC-dimension at most $2^{d+1} - 1$ (see [16]). In particular, the VC-dimension of \mathcal{F} is bounded if and only if the dual VC-dimension is.

Let χ be an m -coloring of the edges of the complete graph K_n with colors q_1, \dots, q_m , and let V be the vertex set of K_n . For $v \in V$ and $i \in [m]$, let $N_{q_i}(v) \subset V$ denote the neighborhood of v with respect to the edges colored with color q_i . We say that χ has *VC-dimension* (or *dual VC-dimension*) d if the set system $\mathcal{F} = \{N_{q_i}(v) : i \in [m], v \in V\}$ has VC-dimension (resp., dual VC-dimension) d .

For $k \geq 3$, $m \geq 2$, and $d \geq 2$, let $r_d(k; m)$ be the smallest integer n such that any m -coloring χ of the edges of K_n with dual VC-dimension at most d contains a monochromatic clique of size k . Even for m -colorings with dual VC-dimension 2, we have $r_2(3; m) = 2^{\Omega(m)}$. Indeed, recursively take two disjoint copies of $K_{2^{m-1}}$, each of which is $(m - 1)$ -colored with dual VC-dimension at most 2 and no monochromatic copy of K_3 . Color all edges between these complete graphs with the m th color, to obtain an m -colored complete graph K_{2^m} with the desired properties. Our main result shows that, apart from a constant factor in the exponent, this construction is tight.

► **Theorem 3.** *For every $k \geq 3$ and $d \geq 2$, there is a constant $c = c(k, d)$ such that $r_d(k; m) \leq 2^{cm}$. In other words, for every m -coloring of the edges of a complete graph of 2^{cm} vertices with dual VC-dimension d , there is a monochromatic complete subgraph of k vertices.*

It follows from the Milnor-Thom theorem [17] (proved 15 years earlier by Petrovskii and Oleinik [18]) that every m -coloring of the $\binom{n}{2}$ pairs induced by n points in \mathbb{R}^d , which is *semi-algebraic* with bounded complexity, has bounded VC-dimension and, hence, bounded dual VC-dimension.

In a recent paper [11], we proved Conjecture 1 for semi-algebraic m -colorings of bounded complexity. Our proof heavily relied on the topology of Euclidean spaces: it was based on the cutting lemma of Chazelle *et al.* [2] and vertical decomposition. These arguments break down in the combinatorial setting, for m -colorings of bounded VC-dimension. In what follows, instead of using “regular” space decompositions with respect to a set of polynomials, our main tool will be a partition result for abstract hypergraphs, which can be easily deduced from the dual of Haussler’s packing lemma [13]. The proof of this partition result will be given in Section 2, while Section 3 contains the proof of Theorem 3.

Overview of the proof. We briefly sketch the idea of the proof of the main theorem. Let χ be an m -coloring of the edges of the complete graph K_n with colors q_1, \dots, q_m , and let V be the vertex set of K_n . Set $\mathcal{F} = \{N_{q_i}(v) : i \in [m], v \in V\}$. Assuming that \mathcal{F} has bounded VC-dimension, we apply a partition lemma to \mathcal{F} discussed in the next section to obtain a vertex partition $V = S_1 \cup \dots \cup S_r$ such that for any pair of vertices $\{u, v\}$ that lies in the same part, very few sets in \mathcal{F} will cross $\{u, v\}$, that is, contain one vertex but not the other. As a consequence, if any part S_t contains a monochromatic K_{k-1} in color q_i , and there is a vertex $v \in S_t$ with large degree with respect to color q_i , then a vertex $u \in N_{q_i}(v)$ can be added to K_{k-1} to produce a monochromatic K_k in color q_i and we are done. Moreover, if there is a “large” part S_t which does not contain a monochromatic K_{k-1} with respect to many of the colors in $\{q_1, \dots, q_m\}$, then we are also done, by induction. If there is no large part with the above property, then the colors of nearly all edges can be “recovered” by much fewer sets in \mathcal{F} . Now we can repeat the argument above on this smaller collection of sets $\mathcal{F}' \subset \mathcal{F}$.

To simplify the presentation, throughout this paper we omit the floor and ceiling signs whenever they are not crucial. All logarithms are in base 2.

2 A partition lemma

Let \mathcal{F} be a set system with dual VC-dimension d and with ground set V . Given two points $u, v \in V$, we say that a set $A \in \mathcal{F}$ *crosses* the pair $\{u, v\}$ if A contains at least one member of $\{u, v\}$, but not both. We say that the set $X \subset V$ is δ -*separated* if for any two points $u, v \in X$, there are at least δ sets in \mathcal{F} that cross the pair $\{u, v\}$. The following *packing lemma* was proved by Haussler in [13].

► **Lemma 4.** *Let \mathcal{F} be a set system on a ground set V such that \mathcal{F} has dual VC-dimension d . If $X \subset V$ is δ -separated, then $|X| \leq c_1(|\mathcal{F}|/\delta)^d$ where $c_1 = c_1(d)$.*

As an application of Lemma 4, we obtain the following partition lemma.

► **Lemma 5.** *Let \mathcal{F} be a set system on a ground set V such that $|V| = n$ and \mathcal{F} has dual VC-dimension d . Then there is a constant $c_2 = c_2(d)$ such that for any δ satisfying $1 \leq \delta \leq |\mathcal{F}|$, there is a partition $V = S_1 \cup \dots \cup S_r$ of V into $r \leq c_2(|\mathcal{F}|/\delta)^d$ parts, each of size at most $\frac{2n}{c_1(|\mathcal{F}|/\delta)^d}$, such that any pair of vertices from the same part S_i is crossed by at most 2δ members of \mathcal{F} . (Here $c_1 = c_1(d)$ is the same constant as in Lemma 4.)*

Proof. Let $X = \{x_1, \dots, x_{r'}\}$ be a maximal subset of V that is δ -separated with respect to \mathcal{F} . By Lemma 4, $|X| = r' \leq c_1(|\mathcal{F}|/\delta)^d$. We define a partition $V = S'_1 \cup \dots \cup S'_{r'}$ of the vertex set such that $x_i \in S'_i$, and for $v \in V \setminus X$, $v \in S'_i$ if i is the smallest index such that the number of sets from \mathcal{F} that cross the pair $\{v, x_i\}$ is at most δ . Such an i always exists since X is maximal. By the triangle inequality, for any two vertices $u, v \in S'_i$, there are at most 2δ sets in \mathcal{F} that cross the pair $\{u, v\}$.

If a part S'_i has size more than $\frac{2n}{c_1(|\mathcal{F}|/\delta)^d}$, we partition S'_i (arbitrarily) into parts of size $\lfloor \frac{2n}{c_1(|\mathcal{F}|/\delta)^d} \rfloor$ and possibly one additional part of size less than $\lfloor \frac{2n}{c_1(|\mathcal{F}|/\delta)^d} \rfloor$. Let $\mathcal{P} : V = S_1 \cup \dots \cup S_r$ be the resulting partition, where $r \leq c_2(|\mathcal{F}|/\delta)^d$ and $c_2 = c_2(d)$. Then \mathcal{P} satisfies the above properties. ◀

3 Proof of Theorem 3

Let d, k_1, \dots, k_m be positive integers. We define the Ramsey number $r_d(k_1, \dots, k_m)$ to be the smallest integer n with the following property. For any m -coloring χ of the edges of K_n with colors $\{q_1, \dots, q_m\}$ such that χ has dual VC-dimension at most d , there is a monochromatic copy of K_{k_i} in color q_i for some $1 \leq i \leq m$. We now prove the following theorem, from which Theorem 3 immediately follows.

► **Theorem 6.** *For fixed integers $d, k \geq 1$, if $k_1, \dots, k_m \leq k$, then $r_d(k_1, \dots, k_m) \leq 2^{cm}$ where $c = c(d, k)$.*

Proof. Let $c = c(d, k)$ be a large constant that will be determined later. We will show that $r_d(k_1, \dots, k_m) \leq 2^c \sum_{i=1}^m k_i$ by induction on $s = \sum_{i=1}^m k_i$. The base case $s \leq k2^{16dk}$ follows by setting c to be sufficiently large.

For the inductive step, assume that $s > k2^{16dk}$ and that the statement holds for all $s' < s$. Thus, we have $m \geq 2^{16dk}$. Set $n = 2^{cs}$ and $Q = \{q_1, \dots, q_m\}$, and let $\chi : E(K_n) \rightarrow Q$ be an m -coloring of the edges of K_n with colors q_1, \dots, q_m such that the set system

$$\mathcal{F} = \{N_{q_i}(v) : v \in V(K_n), q_i \in Q\}$$

has dual VC dimension at most d .

Let $\log^{(j)} m$ denote the j -fold iterated logarithm function, where $\log^{(0)} m = m$ and $\log^{(j)} m = \log(\log^{(j-1)} m)$. For convenience, we will set $\log^{(-1)} m = \infty$. Suppose that there is no color q_i such that χ produces a monochromatic K_{k_i} whose every edge is of color q_i . Otherwise we are done. Then, for every $j \geq 0$ such that $\log^{(j-1)} m > 2^{8dk}$, we recursively construct

1. a set $V_j \subset V$ with $|V_j| \geq n(1 - \frac{1}{\log^{(j-1)} m})$, and
2. an assignment of colors $\chi_j : V_j \rightarrow 2^Q$ to each vertex in V_j with the property that the set system $\mathcal{F}_j = \{N_{q_i}(v) \cap V_j : v \in V_j, q_i \in \chi_j(v)\}$ covers all but at most $8n^2 / \log^{(j-1)} m$ edges of K_n . Here, uv is said to be covered by \mathcal{F}_j if $\chi(uv) = q_i$ implies that $q_i \in \chi_j(u) \cap \chi_j(v)$. Moreover, $|\chi_j(v)| \leq \log^{(j)} m$ for all $v \in V_j$.

We start by setting $V_0 = V$, $\chi_0(v) = Q$ for all $v \in V$, and therefore we have $\mathcal{F}_0 = \mathcal{F}$. Suppose we have V_j, χ_j , and \mathcal{F}_j with the properties described above. Before defining the set V_{j+1} and the assignment of colors $\chi_{j+1} : V_{j+1} \rightarrow 2^Q$, we introduce $B_j \subset E(K_n)$ to be the set of edges that are not covered by \mathcal{F}_j . Hence, $|B_j| \leq \frac{8n^2}{\log^{(j-1)} m}$. We apply Lemma 5 to \mathcal{F}_j , whose ground set is V_j , with parameter $\delta = \frac{|\mathcal{F}_j|}{(\log^{(j)} m)^4}$, and obtain a partition $\mathcal{P} : V_j = S_1 \cup \dots \cup S_r$, where $r \leq c_2(\log^{(j)} m)^{4d}$ and c_2 is defined in Lemma 5, such that \mathcal{P} has the properties described in Lemma 5. For each part $S_t \in \mathcal{P}$, let $Q_t \subset \{q_1, \dots, q_m\}$ be the set of colors such that $q_i \in Q_t$ if there is a vertex $v \in S_t$ such that

$$|\{u \in V_j : \chi(uv) = q_i, uv \notin B_j\}| \geq \frac{n}{(\log^{(j)} m)^2}.$$

Let $Q'_t \subset \{q_1, \dots, q_m\}$ be the set of colors such that $q_i \in Q'_t$ if the vertex set S_t contains a monochromatic copy of K_{k_i-1} in color q_i .

► **Observation 7.** *If there is a color $q_i \in Q_t \cap Q'_t$, then χ produces a monochromatic copy of K_{k_i} in color q_i .*

Proof. Suppose $q_i \in Q_t \cap Q'_t$ and let $X = \{x_1, \dots, x_{k_i-1}\} \subset S_t$ be the vertex set of a monochromatic clique of order $k_i - 1$ in color q_i . Fix $v \in S_t$ such that for $U = \{u \in V_j : \chi(uv) = q_i, uv \notin B_j\}$, we have $|U| \geq \frac{n}{(\log^{(j)} m)^2}$. Notice that if $X \not\subset (N_{q_i}(u) \cap V_j)$, where $u \in U$, then the set $(N_{q_i}(u) \cap V_j)$ crosses the pair $\{x, v\}$ for some $x \in X$. Moreover, $(N_{q_i}(u) \cap V_j) \in \mathcal{F}_j$ since $uv \notin B_j$. Since there are at most $2\delta = \frac{2|\mathcal{F}_j|}{(\log^{(j)} m)^4}$ sets in \mathcal{F}_j that cross $\{x, v\}$, there are at most $\frac{2k|\mathcal{F}_j|}{(\log^{(j)} m)^4}$ sets in $\{N_{q_i}(u) \cap V_j : u \in U\} \subset \mathcal{F}_j$ that do not contain X . On the other hand,

$$|U| - k_i \geq \frac{n}{(\log^{(j)} m)^2} - k_i > \frac{2k|\mathcal{F}_j|}{(\log^{(j)} m)^4},$$

where the last inequality follows from the fact that $|\mathcal{F}_j| \leq n \log^{(j)} m$ and $\log^{(j)} m > 2^{8dk}$. Hence, there must be a neighborhood $(N_{q_i}(u) \cap V_j)$ that contains X , which implies that $X \cup \{u\}$ induces a monochromatic copy of K_{k_i} in color q_i . ◀

By the observation above, we can assume that $Q_t \cap Q'_t = \emptyset$ for every t , since otherwise we would be done.

► **Observation 8.** *If there is a part $S_t \in \mathcal{P}$ such that $|S_t| \geq n / (\log^{(j)} m)^{6d}$ and $|Q_t| \geq \log^{(j+1)} m$, then S_t contains a monochromatic copy of K_{k_i} in color q_i where $q_i \in Q'_t$.*

46:6 Bounded VC-Dimension Implies the Schur-Erdős Conjecture

Proof. For sake of contradiction, suppose $S_t \in \mathcal{P}$ does not contain a monochromatic copy of K_{k_i} in color $q_i \in Q'_t$. Since $Q_t \cap Q'_t = \emptyset$, S_t also does not contain a monochromatic copy of $K_{k_{i-1}}$ in color $q_i \in Q_t$. So if $|Q_t| \geq \log^{(j+1)} m$, we have $|Q'_t| \leq m - \log^{(j+1)} m$. By the induction hypothesis, we have

$$\frac{n}{(\log^{(j)} m)^{6d}} \leq |S_t| < 2^{c(s - \log^{(j+1)} m)}.$$

Since $c = c(d, k)$ is sufficiently large, we have $n < 2^{cs}$ which is a contradiction. \blacktriangleleft

Hence, we can assume that for each part $S_t \in \mathcal{P}$ such that $|S_t| \geq n/(\log^{(j)} m)^{6d}$, we have $|Q_t| < \log^{(j+1)} m$.

We now define the set V_{j+1} and the color assignment χ_{j+1} as follows. Let $V_{j+1} \subset V_j$ be the set of vertices $v \in V_j$ such that v does not lie in a part S_t such that $|S_t| < n/(\log^{(j)} m)^{6d}$. Hence,

$$\begin{aligned} |V_{j+1}| &\geq |V_j| - c_2 (\log^{(j)} m)^{4d} \frac{n}{(\log^{(j)} m)^{6d}} \\ &\geq n - \frac{n}{\log^{(j-1)} m} - \frac{c_2 n}{(\log^{(j)} m)^{2d}} \\ &\geq n - \frac{n}{\log^{(j)} m}. \end{aligned}$$

For each vertex $v \in V_{j+1}$, v lies in a part $S_t \in \mathcal{P}$ with $|S_t| \geq n/(\log^{(j)} m)^{6d}$. We set $\chi_{j+1}(v) = Q_t$, and by the observation above, $|\chi_{j+1}(v)| \leq \log^{(j+1)} m$. Thus, we have the set system $\mathcal{F}_{j+1} = \{N_{q_i}(v) \cap V_{j+1} : v \in V_{j+1}, q_i \in \chi_{j+1}(v)\}$ such that $|\mathcal{F}_{j+1}| \leq n \log^{(j+1)} m$.

Finally, it remains to show that \mathcal{F}_{j+1} covers at least $\binom{n}{2} - \frac{8n^2}{\log^{(j)} m}$ edges of K_n . Let $B_{j+1} \subset E(K_n)$ denote the set of edges that are not covered by \mathcal{F}_{j+1} . If $uv \in B_{j+1}$, then either

1. $uv \in B_j$, or
2. u (or v) lies inside a part $S_t \in \mathcal{P}$ such that $|S_t| \leq \frac{n}{(\log^{(j)} m)^{6d}}$, or
3. both u and v lie inside the same part $S_t \in \mathcal{P}$, or
4. uv is covered by \mathcal{F}_j , but is not covered by \mathcal{F}_{j+1} since $v \in S_t \in \mathcal{P}$ and $\chi(u, v) \notin Q_t$.

By assumption,

$$|B_j| \leq \frac{8n^2}{\log^{(j-1)} m}. \quad (1)$$

The number of edges of the second type is at most

$$\frac{n^2}{(\log^{(j)} m)^{6d}}. \quad (2)$$

The number of edges of the third type is at most

$$\sum_{i=1}^r \binom{|S_i|}{2} \leq c_2 (\log^{(j)} m)^{4d} \left(\frac{2n}{c_1 (\log^{(j)} m)^{4d}} \right)^2 = \frac{4c_2 n^2}{(c_1)^2 (\log^{(j)} m)^{4d}}, \quad (3)$$

where c_1 is defined in Lemma 4. Finally, let us bound the number of edges of the fourth type. Fix $v \in S_t \in \mathcal{P}$ such that $|S_t| > \frac{n}{(\log^{(j)} m)^{6d}}$, and let us consider all edges incident to v that are covered by \mathcal{F}_j . Since v contributed at most $\log^{(j)} m$ sets in \mathcal{F}_j , there are at most $\log^{(j)} m$ distinct colors among these edges. Fix such a color q_i such that $q_i \notin Q_t$, and consider the set of vertices

$$U = \{u \in V_{j+1} : \chi(uv) = q_i, uv \notin B_j\}.$$

By definition of Q_t , we have $|U| < \frac{n}{(\log^{(j)} m)^2}$. Therefore, the number of edges incident to v of the fourth type is at most

$$\log^{(j)} m \frac{n}{(\log^{(j)} m)^2} = \frac{n}{\log^{(j)} m}.$$

Hence, the total number of edges of the fourth type is at most

$$\frac{n^2}{\log^{(j)} m}. \tag{4}$$

Thus by summing (1), (2), (3), (4), and using the fact that $\log^{(j-1)} m > 2^{8dk}$, we have

$$|B_{j+1}| \leq \frac{8n^2}{\log^{(j-1)} m} + \frac{n^2}{(\log^{(j)} m)^{6d}} + \frac{4c_2n^2}{(c_1)^2(\log^{(j)} m)^{4d}} + \frac{n^2}{\log^{(j)} m} < \frac{8n^2}{\log^{(j)} m}.$$

Hence, \mathcal{F}_{j+1} covers at least $\binom{n}{2} - \frac{8n^2}{\log^{(j)} m}$ edges of K_n .

Let w be the minimum integer such that $\log^{(w)} m \leq 2^{8dk}$. Then we have $V_w, \chi_w, \mathcal{F}_w$ with the properties described above. Just as before, let $B_w \subset E(K_n)$ be the set of edges not covered by \mathcal{F}_w . This implies $|B_w| \leq n^2/2^{8dk} < n^2/8$ and $|V_w| \geq 7n/8$. Since

$$\binom{7n/8}{2} - \frac{n^2}{8} \geq \frac{n^2}{4},$$

an averaging argument shows that there is a vertex $v \in V_w$ that is incident to at least $n/2$ edges that are covered by \mathcal{F}_w . Since v contributes at most $\log^{(w)} m < 2^{8dk}$ sets in \mathcal{F}_w , there is a color q_i such that

$$|N_{q_i}(v)| \geq \frac{n}{2 \cdot 2^{8dk}} = \frac{2^{cs}}{2 \cdot 2^{8dk}} \geq 2^{c(s-1)},$$

where the second inequality follows from the fact that $c = c(d, k)$ is sufficiently large. Therefore, by induction, the set $N_{q_i}(v) \subset V$ contains a monochromatic copy of K_{k_ℓ} in color $q_\ell \in \{q_1, \dots, q_m\} \setminus q_i$, in which case we are done, or contains a monochromatic copy of K_{k_i-1} in color q_i . In the latter case, we obtain a monochromatic K_{k_i} in color q_i by including vertex v . This completes the proof of Theorem 6. ◀

4 Concluding remarks

We have established tight bounds for multicolor Ramsey numbers for graphs with bounded VC-dimension. It would be interesting to prove other well-known conjectures in extremal graph theory for graphs and hypergraphs with bounded VC-dimension, especially the notorious Erdős-Hajnal conjecture.

An old result of Erdős and Hajnal [6] states that for every hereditary property P which is not satisfied by all graphs, there exists a constant $\varepsilon(P) > 0$ such that every graph of n vertices with property P has a clique or an independent set of size at least $e^{\varepsilon(P)\sqrt{\log n}}$. They conjectured that this bound can be improved to $n^{\varepsilon(P)}$. Thus, every graph G on n vertices with bounded VC-dimension contains a clique or an independent set of size $e^{\Omega(\sqrt{\log n})}$. In [10], the authors improved this bound to $e^{(\log n)^{1-o(1)}}$. However, the following conjecture remains open.

► **Conjecture 9.** For $d \geq 2$, there exists a constant $\varepsilon(d)$ such that every graph on n vertices with VC-dimension at most d contains a clique or an independent set of size $n^{\varepsilon(d)}$.

References

- 1 N. Alon, J. Pach, R. Pinchasi, R. Radoičić, and M. Sharir. Crossing patterns of semi-algebraic sets. *J. Comb. Theory, Ser. A*, 111:310–326, 2005. doi:10.1016/j.jcta.2004.12.008.
- 2 B. Chazelle, H. Edelsbrunner, L. J. Guibas, and M. Sharir. A singly exponential stratification scheme for real semi-algebraic varieties and its applications. *Theor. Comput. Sci.*, 84:77–105, 1991. doi:10.1016/0304-3975(91)90261-Y.
- 3 F. Chung and R. Graham. *Erdős on Graphs: His Legacy of Unsolved Problems*. A K Peters Series. Taylor & Francis, 1998. URL: https://books.google.com/books?id=Doc_ErUTDcAC.
- 4 F. R. K. Chung. On the ramsey numbers $n(3, 3, \dots, 3; 2)$. *Discret. Math.*, 5:317–321, 1973. doi:10.1016/0012-365X(73)90125-8.
- 5 Paul Erdős. On sequences of integers no one of which divides the product of two others and on some related problems. *Inst. Math. Mech. Univ. Tomsk*, 2:74–82, 1938.
- 6 P. Erdős and A. Hajnal. Ramsey-type theorems. *Discret. Appl. Math.*, 25:37–52, 1989. doi:10.1016/0166-218X(89)90045-0.
- 7 J. Fox, M. Gromov, V. Lafforgue, A. Naor, and J. Pach. Overlap properties of geometric expanders. *Reine Angew. Math. (Crelle's Journal)*, 2012:49–83, 2010. doi:10.1515/CRELLE.2011.157.
- 8 J. Fox, J. Pach, A. Sheffer, A. Suk, and J. Zahl. A semi-algebraic version of Zarankiewicz's problem. *J. Eur. Math. Soc. (JEMS)*, 19:1785–1810, 2014. doi:10.4171/JEMS/705.
- 9 J. Fox, J. Pach, and A. Suk. A polynomial regularity lemma for semialgebraic hypergraphs and its applications in geometry and property testing. *SIAM J. Computing*, 45:2199–2223, 2015. doi:10.1137/15M1007355.
- 10 J. Fox, J. Pach, and A. Suk. Erdős-Hajnal conjecture for graphs with bounded VC-dimension. *Discret. Comput. Geom.*, 61:809–829, 2019. doi:10.1007/s00454-018-0046-5.
- 11 J. Fox, J. Pach, and A. Suk. The Schur-Erdős problem for semi-algebraic colorings. *Israel J. Mathematics*, to appear.
- 12 L. Guth. *Polynomial Methods in Combinatorics*. University Lecture Series. Amer. Math. Soc., 2016.
- 13 D. Haussler. Sphere packing numbers for subsets of the boolean n -cube with bounded Vapnik-Chervonenkis dimension. *J. Comb. Theory, Ser. A*, 69:217–232, 1995.
- 14 D. Haussler and E. Welzl. epsilon-nets and simplex range queries. *Discret. Comput. Geom.*, 2:127–151, 1987. doi:10.1007/BF02187876.
- 15 T. Kóvari, V. Sós, and P. Turán. On a problem of K. Zarankiewicz. *Colloq. Math.*, 3:50–57, 1954. URL: <http://eudml.org/doc/210011>.
- 16 J. Matoušek. *Lectures on discrete geometry*, volume 212 of *Graduate texts in mathematics*. Springer, 2002.
- 17 J. Milnor. On the betti numbers of real varieties. *Proc. Amer. Math. Soc.*, 15:257–280, 1964.
- 18 I. G. Petrovskii and O. A. Oleńnik. On the topology of real algebraic surfaces (russian). *Izv. Akad. Nauk SSSR Ser. Mat.*, 13:389–402, 1949.
- 19 I. Schur. Über die Kongruenz $x^m + y^m = z^m \pmod{p}$. *Jber. Deutsch. Math. Verein.*, 25:114–116, 1917. URL: <http://eudml.org/doc/145475>.
- 20 E. Szemerédi and W. T. Trotter. Extremal problems in discrete geometry. *Combinatorica*, 3:381–392, 1983. doi:10.1007/BF02579194.
- 21 V. N. Vapnik and A. Ya. Chervonenkis. On the uniform convergence of relative frequencies of events to their probabilities. *Theory Probab. Appl.*, 16:264–280, 1971. doi:10.1137/1116025.
- 22 X. Xu, X. Zheng, G. Exoo, and S. P. Radziszowski. Constructive lower bounds on classical multicolor ramsey numbers. *Electr. J. Comb.*, 11, 2004. URL: http://www.combinatorics.org/Volume_11/Abstracts/v11i1r35.html.